



ABN 22 080 933 455

Report ARU-11/002

FINAL REPORT FOR EL 27290 (CONICAL HILL), REYNOLDS RANGE, NORTHERN TERRITORY, AUSTRALIA

By

Kelvin James Hussey
BSc Hons, MAIG

1:250,000 – Napperby SF 53-9

Table of Contents

NORTHERN TERRITORY, AUSTRALIA	1
List of Figures.....	3
List of Tables.....	3
Appendices (Digital data only)	3
INTRODUCTION	4
Background.....	4
Location and access	4
Topography and drainage.....	4
Climate	5
SUMMARY	6
TENURE	7
Mining/Mineral Rights	7
Land Tenure.....	7
Native Title	7
Site Clearances.....	7
Sensitive Areas	8
GEOLOGICAL SETTING	9
Regional Geology	9
Local Geology.....	10
STRATIGRAPHY	10
IGNEOUS ROCKS	13
METAMORPHISM & STRUCTURAL GEOLOGY	14
MINERALISATION	17
PREVIOUS INVESTIGATIONS	18
INVESTIGATIONS BY ARAFURA RESOURCES ON EL 27290	22
RESULTS/DISCUSSION	23
REFERENCES/SOURCES OF INFORMATION	24

List of Figures

- Figure 1. Topographic map showing location EL 27290 and tenement status.
- Figure 2. Approximate location of tenement on NT regional geology map.
- Figure 3. Generalised Reynolds Range Region geology.
- Figure 4. Distribution of Stafford Event metamorphism.
- Figure 5. Distribution of Chewings Event metamorphism.
- Figure 6. Distribution of Alice Springs Orogeny metamorphism.
- Figure 7. Geological map showing location of EL 27290.
- Figure 8. Detailed geological map of EL 27290.

List of Tables

- Table 1. Summary of historic exploration activity.

Appendices (Digital data only)

- Appendix 1. Rock chip and assay data.

INTRODUCTION

Background

EL 27290 is part of a number of tenements in the Aileron-Reynolds Range area that Arafura Resources has the mineral exploration rights to. These tenements include EL 23571 (Reynolds Range), SEL 23671 (Aileron), EL 24548 (Yalyirimbi Range), EL 24741 (Woodforde) and EL 27291(Boundary Bore).

The Reynolds Range area is prospective for numerous styles of mineralisation with U, Au, As, Sb, Ag, Fe, Sn, Ta, W, Mo, Cu, Pb, Zn, Ni, REE, P, Th and talc occurrences known in the region. Of these, Arafura Resources is principally interested in exploring for economic REE mineralisation.

Several companies conducted exploration activities in the adjacent SEL 23671, and failed to identify or realise the potential of a globally significant REE deposit that was partly outcropping at Nolans Bore. Resource drilling by Arafura Resources to date, confirms the Nolans Bore deposit has a total resource of 30.3 Mt @ 2.8% REO, 12.9 % P₂O₅ and 0.44 lb/t U₃O₈ [see Goulevitch (2008) for details]. This is considered encouraging because the Nolans Bore- type REE mineralisation may occur elsewhere in the region.

Elevated levels of phosphate-hosted rare earth elements (REEs) mineralisation was discovered in the Nolan's Bore area by PNC Exploration (Australia) Pty Ltd in 1995 (Thevissen, 1995). This occurred during follow-up of an airborne radiometric anomaly as part of that company's uranium exploration program along the Reynolds Range. Total REE levels of 5-7% were reported from selected grab samples of apatite which crop out sporadically within an area some 1000 x 800 metres in extent around the Bore. A distinct ground-radiometric response correlates closely with this area of apatite development.

Location and access

EL 27290 (Conical Hill) is located about 25 kilometres west-northwest of Aileron in the central-southern part of the Northern Territory. The Stuart Highway is the main north-south arterial highway through the Northern Territory passing the Aileron Roadhouse about 130 kilometres north of Alice Springs.

The tenement can be reached via a seldom used station track traversing the southern side of the ranges from the Stuart Highway via Old Alb's bore and passing through the tenement on the way to Boundary Bore. There are no other vehicle tracks in the tenement area.

Topography and drainage

The Reynolds Range is a relatively narrow (5-10 kilometre wide, 90 kilometres long), northwest trending belt of steep hills with deeply incised drainages. Elevations in the Reynolds Range exceed 750 metres above sea level, with the highest peaks reaching over 1000 metres above sea level (eg. Mt Thomas, 1116 metres). Mt Freeling (1005 metres) and Mt Dunkin (930 metres) are other prominent landmarks in the region.

Mt Freeling is part of the ranges and lies immediately north of EL 27290. The topography on EL 27290 flanks the ranges and is much more subdued compared to the main range of hills. Most of the tenement gently slopes towards the south with drainage lines shedding off the ranges.

Climate

The climate is characterised by long hot summers and short mild winters. Temperatures regularly exceed 40°C in summer with rare frosts in winter. The average rainfall is about 280 mm, most of which falls between October and March, but both frequency and amount are erratic.

The Aileron-Reynolds project area is part of the Burt Plain Bioregion as defined by Connors (2004). Connors (2004) indicates that the Ranges are broadly classified as “*mixed species low open woodlands*” which is described as spinifex hummocky grassland with a mixed low open-woodland overstory (principally Eucalyptus and Hakea species). The southern slope of most rocky ridges also contains characteristic isolated stands of native Pines. The open valleys between the Ranges are classified as “*tall open shrublands*” which is principally *Acacia kempeana* (Witchetty Bush) tall open-shrubland with Cassia and Eremophila open-shrubland understory, but also includes an unmapped area of *Acacia aneura* (Mulga) tall open-shrubland with Cassia and Eremophila open-shrubland understory. Extensive *Acacia aneura* (Mulga) tall open-shrubland dominates the southern sides on the ranges. Major drainages contain good stands of River Red Gums, Bloodwood and lesser Beantree.

SUMMARY

EL 27290 was granted to Arafura Resources on 2 December 2009 and relinquished on 18 October 2010. This report documents all exploration activity by Arafura Resources on EL 27290 during the period of grant.

EL 27290 was only held for less than one year and consequently limited exploration activities have occurred within the tenement area.

Office-based desktop studies including literatures searches and GIS studies were conducted in 2009-2010 using NTGS regional and company datasets. These studies identified a number of geological targets to be investigated. These targets corresponded to mapped calcsilicate rocks, currently thought to be part of the Reynolds Range Group. Some exploration targets were investigated in late 2009 and others in early 2010. However the extreme wet conditions of 2010 prevented ongoing access to the tenement and the short term of tenure meant that only limited on-ground exploration could be conducted during the period of grant.

Arafura visited a total of 6 sites in 2009, while under application, as it was working in the area and a further 2 sites were visited during the period of grant. Arafura attempted additional preliminary field reconnaissance of EL 27290 on three occasions between June and August 2010. However all attempts at vehicular access during mid-2010 proved fruitless, largely due to the extremely wet boggy conditions and flowing creeks on the southern side of the Reynolds Range.

The sites visited by Arafura contained no economic mineralisation of interest. Despite this Arafura identified a potential for Sn-P-Y-HREE skarn- or vein-type mineralisation in the region and more systematic follow up exploration is warranted.

Nolans Bore-type mineralisation was not identified in EL 27290 however the absence of detailed airborne geophysical data means that the potential for Nolans Bore-type mineralisation has not been thoroughly evaluated.

TENURE

Mining/Mineral Rights

Exploration Licence 27290 (Conical Hill) was 100% held by Arafura Resources Limited (ACN 080 993 455) under Miners Right number 13249. Arafura Resources holds all Mineral Rights.

The application of 3 sub-blocks (about 10 square kilometres) was granted to Arafura Resources on 2 December, 2009 for a period of 4 years (Figure 1). EL 27290 was surrendered on 18 October 2010 and the area incorporated as part of a new larger EL application (EL 28498).

Land Tenure

The land tenure under EL 27290 (Conical Hill) is PPL 1097 (Aileron).

- Aileron Station, PPL 1097 – NT parcel 00703 is owned by Mr Garry Dann of Aileron Station (Waite River Holdings Pty Ltd), (phone 08 8956 9705, fax 08 8956 8535).

Native Title

Arafura Resources has negotiated and executed an Exploration Agreement with the Central Land Council (on behalf of registered Native Title Claimants).

In December, 2003, an Introductory Meeting was held at Nolan's Bore with members of the relevant Native Title groups. A further meeting with CLC officers and relevant Native Title groups was held at Aileron on 31 March, 2006, where the adjacent EL 23571 and SEL 23671 were confirmed as part of the Exploration Agreement.

EL 27290 was not subject to this original Exploration Agreement but it was Arafura's intention to bring the tenement into the Agreement once a suitable meeting was arranged. No such meetings were held during the grant period. Despite this the area was still appropriately cleared for exploration activity as a part of a regional clearance while under application in late 2009. As a result, there are no Native Title impediments to continued exploration other than holding appropriate consultations, avoiding activity on identified sacred sites and paying agreed amounts of financial compensation.

The terms of the Exploration Agreement provide for continuation of exploration on the area of the proposed mining tenement while the mining agreement is being negotiated with the registered Native Title Claimants.

Site Clearances

Under the terms of the Exploration Agreement, Arafura must provide all relevant details of its proposed exploration activities to be conducted. The CLC must advise if clearances are necessary and then, if required, conduct clearances and provide details of exclusion zones as advised by the Native Title holders. Under the Exploration Agreement, the CLC is required to provide all necessary Sacred Site Clearances and details of the exclusion zones to allow exploration activities to progress in a timely manner.

Arafura submitted a request to the CLC for reconnaissance access to the surrounding areas in 2007/08. Due to the nature of the area, the CLC advised that access could not be given without substantial clearances. Arafura therefore decided to acquire hyperspectral survey data over the entire

region in 2008 as this technique could potentially serve as a method to refine exploration targets and hence reduce the amount of clearance required.

Arafura provided all relevant details of its proposed reconnaissance exploration activities over the greater Aileron-Reynolds project area to the CLC in July 2009. On 25 September, 2009, the CLC provided Sacred Site Clearance Certificate 2009-075 detailing all exclusion zones within the outlined exploration area. This clearance included the area encompassed by EL 27290.

Sensitive Areas

There are no documented sensitive areas in the tenement area.

GEOLOGICAL SETTING

Regional Geology

The Arunta Region contains more than 200 000 km² of metamorphic rocks in the southern parts of the NT and has been subdivided into three distinct geological regions by the NTGS, the Aileron, Warumpi and Irindina Provinces (Figure 2). The Arunta Region is unconformably overlain by sediments of the Neoproterozoic to mid-Palaeozoic Ngalia, Georgina, Amadeus and Wiso Basins.

The Aileron Province predominantly consists of Palaeoproterozoic sedimentary and igneous rocks that have undergone greenschist to granulite facies metamorphism. The majority of the preserved metasedimentary and igneous rock units in this region were deposited or emplaced prior to the 1740-1690 Ma Strangways Orogeny (e.g. Scrimgeour 2003, Hussey *et al* 2005, Clauoué-Long *et al* 2008a, 2008b). This event appears to have affected the entire Aileron Province to some degree, as opposed to the Mesoproterozoic 1595-1570 Ma Chewings Event that appears to be localised within the central and southern(?) parts of Aileron Province (e.g. Hand and Buick, 2001, Fraser, 2004). The 1800 Ma Stafford and 1790-1770 Ma Yambah Events also appear to be present throughout the Aileron Province, with extensive bimodal igneous activity, associated sedimentation and localised Low Pressure-High Temperature metamorphism.

Most of the exposed Aileron Province was metamorphosed to greenschist or lower amphibolite facies conditions during the 1740-1690 Ma Strangways Orogeny, with an apparent localised abundance of 1810-1700 Ma igneous activity and deformation in parts. The central-southern parts of the Aileron Province preserves an east-west zone of granulite facies metamorphic rocks associated with the Strangways Orogeny. Regions of the Aileron Province have also been subject to younger (1640-1500 Ma) periods of magmatism and localised metamorphism.

Current views on the depositional and tectonic setting of the Aileron Province are based on recent geochemical, isotopic and igneous studies and the contained mineral systems. These favour a rifted continental crust or evolving backarc setting in the early parts of the depositional history [e.g. Hussey *et al* 2005, Hoatson *et al* 2005 Matthew Cobb (PhD student, Curtin University) *pers. comm.*, 2005], with a prolonged tectonothermal convergent event in the Strangways Orogeny. Hussey *et al.* (2005) and Hoatson *et al.* (2005) argue for contiguous sedimentation and bimodal igneous activity during Stafford Event. This Event is thought to be responsible for the development of localised(?) deep-marine basins in the Arunta Region, as opposed to contemporaneous subaerial to shallow-water volcanism and sedimentation in the adjacent Davenport Province.

The Aileron Province contains temporal equivalents of the gold-bearing Granites-Tanami and Tennant Creek Regions and regional aeromagnetic data suggest lateral continuity between these Regions. The Aileron Province is therefore regarded as part of the North Australian Craton, however, localised facies variations and differences in sedimentary environments are evident (e.g. Hussey *et al* 2005).

The Warumpi Province in the south and southeast of the Arunta Region (Figure 2) contains a younger package of metasedimentary and volcanic rock types with protoliths in the range 1690-1600 Ma (Scrimgeour *et al* 2003). The Province was variably metamorphosed in the 1640 Ma Leibig Orogeny, 1570 Ma Chewings and the 1150 Ma Teapot Events.

Unmetamorphosed Neoproterozoic to Palaeozoic marine and terrestrial sedimentary rocks of the Georgina, Ngalia and Amadeus Basins surround and unconformably overly the Arunta Region. Contemporaneous Neoproterozoic to Cambrian strata of the Harts Range Group (Buick *et al* 2001, Maidment *et al* 2004, Buick *et al* 2005) are also caught up within the eastern parts of the Arunta Region in the newly defined Irindina Province (Scrimgeour, 2003). This revision and reinterpretation of the Arunta Region has significant geological implications and has come about largely as a result of several extensive chronological, metamorphic and metallogenic studies in the eastern Arunta Region (eg Miller *et al* 1998, Mawby *et al* 1998, 1999, Hand *et al* 1999a, b, Buick *et al* 2001, Scrimgeour and Raith, 2001, Hussey 2003, Maidment *et al* 2004, Buick *et al* 2005, Clauoué-Long and Hoatson, 2005, Close *et al* 2005, Hussey *et al* 2005).

Geochronological and metamorphic studies have shown that the rocks of the Harts Range Group in the Irindina Province are variably metamorphosed to transitional granulite facies in the (480-450 Ma) Ordovician Larapinta Event. This high-grade event is followed by lower-grade Devonian to Carboniferous deformation and granite and pegmatite intrusion. Interestingly, the high-grade Larapinta Event appears to have had little influence on the thermal history of the surrounding rocks of the Aileron Province, and apart from rare exceptions appears to be largely restricted to the Irindina Province (Maidment 2004, Close *et al* 2005, Hussey *et al* 2005, Claoué-Long and Hoatson, 2005).

Many of the fault bounded contacts between the various units within the Arunta and surrounding regions are attributed to the (390-300 Ma) Devonian-Carboniferous Alice Springs Orogeny. Most of the fault movements within the adjacent Georgina Basin also appear to be related to the Ordovician Larapinta Event and Devonian-Carboniferous Alice Springs Orogeny.

Localised carbonatite occurs at Mud Tank (730 Ma), Mt Bleachmore and also in the Casey Inlier area in the central and southern parts of the Aileron province. The carbonatite ages the latter two regions are unknown but it is conceivable that both are about 730 Ma. A small potassic alkaline igneous complex, the Mordor Igneous Complex that has lamphyrophyric affinities (Barnes *et al* 2008) was emplaced in the southern-central parts of the Aileron Province at 1132 Ma (Claoué-Long & Hoatson, 2005).

Local Geology

(Modified after McGilvray 2006)

STRATIGRAPHY

Palaeoproterozoic

The Lander Rock beds are the oldest known outcropping rocks in the area. The Lander Rock package is a suite of dominantly quartzose and pelitic sediments with a facies transition in the northwest, to alternating pelites and psammites, in the Mt Stafford Beds. At least some parts of the Lander Rock beds preserve sedimentary structures (Bouma sequences) indicative of sedimentation below storm wave base. Major outcrops occur in the Lander River Valley north of the Reynolds Range and in the vicinity of Harverson Pass (Figure 3). The metamorphic grade varies from lower greenschist facies in the northwest of the Reynolds Range to granulite facies in the southeast. Minor sills or dykes of mafic rocks occur in the package. The timing of deposition of the Lander Rock Package is unclear although 1795-1806 Ma granite intrusives (Worden *et al* 2008) and U-Pb SHRIMP detrital zircon ages provide a rough maximum estimate of 1806-1840 Ma (Vry *et al* 1996, Claoué-Long 2003, Claoué-Long *et al* 2005, Claoué-Long *et al* 2008a). It is important to note that recent dating suggests the Lander Rocks beds can be divided into at least two stratigraphic units based on zircon provenance patterns and the presence of a younger zircon population in some areas (Claoué-Long 2003, Claoué-Long *et al* 2005, Claoué-Long *et al* 2008a).




Figure 3: Generalised geology of the Reynolds Range Region (modified after Stewart, 1981). Magnetic data indicates that the bulk of the regions covered by recent alluvium are underlain by granite/granitic gneiss (from Hand & Buick 2001).

The Reynolds Range Group is sub-divided into four stratigraphic units (Buick *et al.* 1999). The basal Quartzite Unit, the Mt. Thomas Quartzite, is a mature orthoquartzite that unconformably overlies the Lander Rock Package in the northwest of the Reynolds Range (Figure 3). The unit varies in thickness from ~200 metres to 550 metres cropping out along the length of the range. The lower units are predominantly conglomeratic with minor pebbly arkose rocks. The upper intervals are pelitic and generally ferruginous. A lateral facies change occurs from the northeast to the southwest across the range from basal conglomerates into homogenous pelitic rocks. Relict sedimentary structures indicate a high-energy, intertidal depositional environment (Buick *et al.* 1999).

The Lower Calcsilicate Unit forms the basal unit of the group in the southern margin of the Reynolds Range. This unit can be age constrained as an equivalent to the Mt Thomas Quartzite and by the intruding Napperby Gneiss (metagranitoid). The unit is composed of finely layered, carbonate-poor calcsilicate rocks rich in clinopyroxene, plagioclase and grossular-andradite garnet locally interlayered with white quartzites and rare marbles. The unit is strongly metamorphosed and intensely deformed lacking sedimentary structures (Buick *et al.* 1999).

The Pelite Unit which was previously part of the Pine Hill Formation achieves a minimum thickness of 500 metres to 600 metres. Pelitic rocks are interlayered with thin sheets of fine grained siltstone and sandstone interpreted as storm deposits (Buick *et al.* 1999).

The Upper Calcsilicate Unit encompasses the previously defined Algamba Dolomite Member and the Woodforde River Beds. The unit achieves a maximum thickness of about 250 metres to 300 metres along the length of the Reynolds Range except in the central part where the maximum thickness is only 20 metres. The unit occurs as a series of lenses within the Pelite Unit dominated by interlayered limestone and dolomite locally intercalated with pelites and psammities. Stromatolites and sedimentary structures, *i.e.* climbing ripples, are preserved where rocks are metamorphosed at a regional low grade (Buick *et al.* 1999).

Neoproterozoic

Ngalia Basin rocks were deposited between the Neoproterozoic to the Late Carboniferous (Wells & Moss, 1983). The rocks are an important component of the adjacent EL 24548, but do not occur in EL 23571.

Tertiary

Cainozoic sediments occur in sedimentary basins outside of EL 23571 and will not be discussed in this report.

Geological research in Tertiary basins (Senior *et al* 1995) has defined three weathering events which affected Arunta igneous and metamorphic basement rocks and lacustrine and fluvial Tertiary sedimentary rocks. The weathering events will be discussed herein.

Weathering Event A occurred during the Late Cretaceous to Early Tertiary (Palaeocene). A trizonal profile was developed in basement rocks over a widespread area of the Arunta Region and at the base of surrounding Tertiary Basins. The trizonal profile consists of a basal kaolinitic zone up to 10 metres thick that grades into a multicoloured mottled zone up to 10 metres thick. The mottled zone is overlain by a ferruginous zone up to 8 metres thick (Senior *et al* 1995). The weathering profile is developed in basement rocks and the Mesozoic Hooray Sandstone, and is overlain by Palaeocene sediments in Tertiary basins.

Weathering Event B affects the upper part of the Ambalindum Sandstone Member immediately beneath the Delaney Mudstone Member in the Hale Basin, located in the eastern part of the NTGS Alice Springs 1:250,000 Geology Map Sheet. The upper part of the Ambalindum Sandstone Member is friable and yellow, having a mottled appearance in parts. The weathering event occurred prior to the Middle Eocene. Little evidence exists outside of the Hale Basin for this weathering event (Senior *et al* 1995).

Weathering Event C affects the upper part of the Tug Sandstone Member of the Hale Formation in the Hale Basin. The weathering event preceded deposition of the Waite Formation in the Waite Basin, or equivalents of the Waite Formation.

Quaternary

Further uplift in the Reynolds Region, and northern Arunta Region has resulted in deposition of red earth and alluvium from uplifted areas and continued movement of colluvium down present-day hillslopes. Calcrete has precipitated along stream channels, evaporites have formed in playa lakes, and sand plains and Aeolian dunes have developed in low lying areas (Stewart, 1981).

IGNEOUS ROCKS

Palaeoproterozoic

Based on recent high precision SHRIMP U-Pb dating of zircons in igneous rocks by the NTGS (Worden *et al* 2008), granitic rocks of the Reynolds Range region can be subdivided into two age-related suites. The existence of two igneous suites requires:

1. emplacement of the first granitic suite into the Lander Rock package,
2. uplift and erosion,
3. deposition of the Reynolds Range Group, and
4. emplacement of the second igneous suite

The emplacement age of the first igneous suite is now well constrained at about 1795-1805 Ma (Worden *et al* 2008). This suite crops out on the northern side of the Reynolds Range, is contemporaneous with LP/HT metamorphism and partial melts at Mount Stafford (the Stafford Event) and provides localised evidence for bimodal magmatism, exemplified by the presence of contemporaneous mafic igneous rocks occur in the Anmatjira Orthogneiss.

From southeast to northwest, the first granitic suite includes the Boothby Orthogneiss (1806 ± 4 Ma, Worden *et al* 2008), Yaningidjara Orthogneiss [1798 ± 4 Ma, Worden *et al* (2008) which is within error of the 1806 ± 6 Ma age by Vry *et al* (1996)], Mount Airy Orthogneiss (1799 ± 3 Ma, Worden *et al* 2008), Harverson Granite (1799 ± 3 Ma, Worden *et al* 2008), Anmatjira Orthogneiss [1798 ± 3 Ma by Worden *et al* 2008 and 1802 ± 3 Ma by Rubatto *et al* (2006) about 15 kilometres north of the first location near Mount Stafford]. Early SHRIMP U-Pb determinations on zircon from these igneous rocks by Collins & Williams (1995) are much less precise and are discounted in favour of more recent CL-assisted SHRIMP U-Pb dating by Worden *et al* (2008) and Rubatto *et al* (2006). Rubatto *et al* (2006) also determined that the LP/HT metamorphism at Mount Stafford occurred between ~1795 and 1805 Ma.

The second slightly younger igneous suite appears to be about 1770-1785 Ma and probably reflects the Yambah Event in this region. The suite mainly outcrops the Reynolds Range and further south, although the Possum Creek Charnokite [1774 ± 6 Ma, Collins and Williams (1995)] and the Tyson Creek Granulites in the Anmatjira Range are similar age. The age of second igneous suite is typically poorly constrained with larger errors, particularly those in the Reynolds Ranges. These units are all high level granites that contain metasedimentary enclaves and have a peraluminous geochemical signature. The assimilation of sedimentary units causes significant zircon inheritance issues making interpretation of magmatic zircon ages difficult (eg Smith 2001). The differentiation of some granitic units is unclear based on current published maps and careful remapping is needed.

The second igneous suite includes the Warimbi Schist [1785 ± 22 Ma, Collins & Williams (1995)], Coniston Schist [1780 ± 10 Ma, Smith (2001)] and Napperby Gneiss [1780 ± 10 Ma, Collins & Williams (1995)]. The Yakalibadgi Microgranite probably also belongs in this suite as do a number of undifferentiated granites and gneisses that intrude the Reynolds Range Group (see Stewart and Pillinger 1981).

METAMORPHISM & STRUCTURAL GEOLOGY

The Arunta Region was shaped by two major intervals of tectonism. The first major tectonic interval occurred during the Palaeo- to Mesoproterozoic, 1850-1560 Ma, and was associated with multiple episodes of regional medium to high temperature metamorphism and magmatism (Hand & Buick, 2001). The second major tectonic interval occurred in the early to mid-Palaeozoic, about 490 to 300 Ma, and was associated with north-south intraplate extension and subsequent north-south convergent deformation (Hand & Buick, 2001). Regional structures produced during each period of tectonism in the Reynolds Range Region are discussed by Hand & Buick (2001).

The first tectonic interval is defined by three main tectonic events, the 1805-1795 Ma Stafford Event, the 1785-1770 Ma Yambah Event, and the 1595-1560 Ma. Chewings Orogeny. There has been significant debate about the Strangways Orogeny in the Reynolds and Anmatjira Ranges. Historically the Strangways Orogeny was about 1780-1720 Ma however recent revision by the NTGS identifies the Yambah Event (1785-1770 Ma) and the Stangways Orogeny (about 1740-1690 Ma). All published literature still refers to the historic usage of Strangways Orogeny. The Yambah Event occurs in the Reynolds Range region and the affect of the Strangways Orogeny as newly defined needs to be resolved.

The Stafford Event is based on LP/HT metamorphism and igneous relationships in the Mount Stafford area. The first igneous suite noted above is coincident with the Stafford Event and includes the Harverson Granite highlighted in Figure 4. The Lander Rock Package around the Harverson Granite is characterised by the growth of andalusite and cordierite (Dirks *et al* 1991; Vry & Cartwright, 1998). The contact metamorphic porphyroblasts overprint a biotite-quartz-muscovite foliation which indicates prior regional deformation to granite emplacement. In other parts of the northwest Reynolds Range, muscovite±biotite bearing greenschist assemblages, (Dirks *et al* 1991), define a sub-vertical northwest-southeast trending foliation, (Stewart, 1981; Dirks & Wilson, 1990). Fold structures are truncated to the overlying Reynolds Range Group in an unconformity that dies to the southeast, indicating the Stafford Tectonic Event may have been localised in the northwestern part of the Reynolds Range, and Anmatjira Range, region (Hand & Buick, 2001).

Contact metamorphic assemblages formed in the Reynolds Range Group around the granitic precursors of the Warimbi and Coniston Schists during intrusion of the second igneous suite at around 1785-1770 Ma (Collins & Williams 1995). Contact aureoles in meta-pelites adjacent to the Warimbi Scist are andalusite and cordierite bearing. The stability of these assemblages indicate maximum P-T conditions of 550°C and 3.5 kilobars (Xu *et al* 1994; Mahar *et al* 1997). Scapolite porphyroblasts in anorthite-bearing marbles adjacent to the Coniston Schist also give maximum temperatures of 550°C (Buick & Cartwright, 1994). Contact metamorphic blasts surrounding the Warimbi Schist contain straight or gently curved internal foliations defined by muscovite-quartz±biotite. Curved inclusion trails indicate the growth of the contact metamorphic assemblages occurred during deformation. Inclusion trails are reported to show systematic changes in orientation defining gentle folds. The orientation of folding is not clear but has been postulated as a southeast trending foliation based on findings in other parts of the Arunta Inlier (Hand & Buick, 2001; Goscombe, 1991; Collins & Sawyer, 1996).

The Chewings Orogeny produced a nearly continuous northeast-southwest transition in metamorphic grade from greenschist to granulite facies along the length of the Reynolds Range (Figure 5). Meta-pelitic rocks of the Reynolds Range Group are transformed from phyllites to andalusite±cordierite-bearing schists to migmatitic granulites (Dirks *et al.*, 1991; Hand & Dirks, 1992; Williams *et al* 1996 & Buick *et al* 1998). The metamorphic field gradient is summarised by Hand and Buick (2001) by the metamorphic zones: muscovite-chlorite±biotite; texturally stable Strangways Orogeny andalusite and cordierite; first appearance of sillimanite and; stable co-existence of cordierite-spinel assemblages.




Figure 4: Regional distribution of metamorphism inferred to be associated with the Stafford Event (from Hand & Buick, 2001).




Figure 5: (a) Simplified geological map of the Reynolds Range Region showing the metamorphic zones associated with the 1595-1570 Ma Chewings Event. (b) Metamorphic zones in the Reynolds Range defined by the assemblages produced during partial melting. Assemblages in the near minimum melting zone include ilmenite-magnetite-bearing leucosomes in migmatized granite. In the granulite zone, leucosomes contain cordierite ± garnet ± orthopyroxene (from Hand & Buick, 2001).

The higher grade regions of the Reynolds Range are further sub-divided by Hand and Buick (2001) based on the leucosome assemblages that formed during partial melting. Upper amphibolite regions show immediate upgrade of the sillimanite isograd, and volumetrically minor leucosomes are pegmatitic in character with simple mineralogies that reflect water-saturated melt (Buick *et al* 1998). At slightly higher grades leucosomes contain ilmenite-magnetite intergrowths that form via breakdown of biotite (Hand & Dirks, 1992). The highest grade granulite leucosomes contain cordierite and/or garnet or orthopyroxene and formed during fluid-absent dehydration reactions that consumed biotite and sillimanite.

Partial melting assemblages overprint the gneissose layering suggesting high temperature metamorphism outlasted pervasive deformation (Hand & Buick, 2001). Granulite and upper amphibolite assemblages are aligned parallel to the axial surface of the regional, upright, southeast-trending, isoclinal folds (Hand & Buick, 2001). The upright folds reflect around 50% shortening and can be traced along the length of the Reynolds Range (Dirks & Wilson, 1990). Many of the macro-scale folds within northwest-southeast regional folds are doubly plunging, (Stewart *et al* 1980; Dirks & Wilson, 1990), which represents significant vertical extension (Hand & Buick, 2001). In the lower grade northwestern Reynolds Range, the axial surface fabric overprints approximately 1785 Ma contact metamorphic minerals.

In the Reynolds Range, the regional fabric has been deformed on all scales by conjugate, steeply-dipping shear and crenulation bands that, in geometry, represent conjugate kink bands (Dirks & Wilson, 1990; Hand & Dirks, 1992). The dominant kink set trends approximately east-west plunging between 0° and 70° east. The subordinate kink set trends approximately north-south and plunges to the north (Hand & Buick, 2001). Zircons from leucosomes within the crenulation bands have been aged at 1570 Ma (Hand *et al* 1995; Williams *et al* 1996), which confirms development of structures during the Chewings Orogeny (Hand & Buick, 2001).

Proterozoic structures in the Reynolds Range are heavily dissected by southeast and east trending shear zones associated with the 400-300 Ma Alice Springs Orogeny (Hand & Buick, 2001). Micaceous greenschist to lower amphibolite assemblages are dated to 330-300 Ma (Cartwright *et al* 1999).

Collins and Teyssier (1989), interpret the overall geometry of the Reynolds-Anmatjira Ranges to have formed in a transpressional setting with a northeast-plunging lineation representing a component of sinistral movement during the Alice Springs Orogeny, resulting in juxtaposition of granulites against lower grade rocks in the southwestern Reynolds Range (Dirks *et al* 1991).

The metamorphic grade of Alice Springs Orogeny structures increases to the southwest (Figure 6) such that shear zones in the southwest of the Reynolds Range contain kyanite, staurolite and sillimanite-bearing assemblages in metapelite, (Dirks *et al* 1991) with P-T conditions of 5-5.5 kilobars and 550-600°C. In the southeastern Anmatjira Ranges, the shear zones contain andalusite and staurolite assemblages in meta-pelite, with P-T conditions of 4 kilobars and 580°C (Xu *et al* 1994). In the central and northwest Reynolds Range the shear zones are associated with greenschist or lower-grade metamorphism (Dirks *et al* 1991). Accompanying the increase in metamorphic grade is an increase in the number and width of the shear zones, with zones in the southeastern Reynolds Range up to 300m wide (Hand & Buick 2001).

Episodic mild uplift and warping consisting of limited upward doming of ranges and minor tilting continued through the Palaeozoic and Cainozoic to present day (Senior *et al* 1995).




Figure 6: Metamorphic zones defined by mid-Palaeozoic metapelitic shear zone assemblages in the Reynolds Range Region (from Hand & Buick 2001).

MINERALISATION

Relevant company reports and descriptions of the Reynolds Range region by the NTGS describe numerous occurrences of mineralisation. These include copper-lead-zinc, gold, tungsten, tin, tantalum, rare earth elements, mica, nickel, chromium, semi-precious stones, talc, iron and uranium. A variety of mineralisation styles have potential in the Reynolds Range region but few mineralisation styles have identified resources that would be considered prospective for economic development.

The Nolans Bore Phosphate- Rare Earth Element- Uranium deposit is considered to be the most prospective economic development in the area. Nolans Bore occurs in the adjacent SEL 23671 and is a globally significant REE resource containing 30.3 Mt @ 12.9% P₂O₅, 2.8% REO and 0.44 lbs/T (202 ppm) U₃O₈ (Goulevitch 2008) and is currently undergoing a Bankable Feasibility Study.

PREVIOUS INVESTIGATIONS

EXPLORATION PRIOR TO 1996

(modified after Drummond 2003)

The following historic exploration summaries may not be explicitly associated with EL 27290 but are relevant when considering regional prospectivity of the Aileron-Reynolds project area. Reference is made to historic EL 9672 (Dragons Lair) and contemporary ELs 23671 (Aileron), 23571 (Reynolds Range), 24548 (Yalyirambi Range) and 24741 (Woodforde) in the following exploration summaries. The relevant historic exploration licences and activities are summarised in Table 1 and are detailed further below.

Table 1: Summary of historic exploration.

Years	Tenement(s)	Exploration Company	Exploration Targets/Commodities	NT Department of Resources Open File Company Report(s)
1977-1978	EL 1294	CSR Minerals & Chemical Division	Base metals, tungsten and uranium.	CR1979-0198
1977-1979	EL 1444	Otter Exploration	Base metals, tin-tungsten, uranium and REE.	CR 1979-0021, CR 1980-0056, CR 1980-0252.
1981-1983	EL 2942, EL 3075, EL 3084, EL 3088.	BHP Exploration	Diamonds, base metals.	CR 1983-0015, CR 1983-0289.
1982-1983	EL 3506	J Weir	Uranium and REE	CR 1983-0216
1983-1984	EL 4188	BHP Exploration	Possible carbonatite	CR 1984-0117
1987-1990	EL 5511	Colchis Mining Corporation	Gold and base metals	CR 1989-0020, CR 1990-0036.
1988-1989	EL 5901	Track Minerals	Gold and base metals	CR 1989-0704
1991-1993	EL 7344, EL 73445	Poseidon Gold	Gold and base metals	CR 1993-0410
1993-2001	EL 7343	Poseidon Gold & Exodus Minerals	Gold and base metals	CR 1999-0255 CR 2001-0270
1993-1994	EL 8117	Tidegate	Gold	CR 1994-0589
1994-1996	EL 8411	PNC Exploration	Uranium	CR 1995-0266, CR 1996-0187
1995-1998	EL 9146	Aberfoyle Resources	Gold	CR 1996-0692, CR 1997-0688
1996-1998	EL 9672	Homestake Gold of Australia	Gold and iron	

CSR minerals (1977-1978) EL 1294.

This historic licence covered most of EL 23571 and the north central portion of the adjacent EL 24548. The licence area was considered prospective for base metals, tungsten and uranium. An airborne radiometric survey was flown in 1977. Thirty seven airborne radiometric anomalies were checked along with stream sediment sampling and rock chip sampling.

Reconnaissance sampling located elevated rock chip values in the range 190 to 1620m ppm U₃O₈. These were obtained from areas of secondary uranium mineralisation with the highest values outside EL 23571 in the Yalyirimbi Range. The best rock chip assay within EL 23571 was 160 ppm U₃O₈. One of these sites does not correspond with a significant airborne radiometric anomaly and its plotted location is suspect. No Sn-W or base metal geochemistry of significance was identified. None of the anomalies were considered prospective.

Otter Exploration (1977-1979) EL 1444.

EL 1444 was located north of EL 23571 and covered the northern portion of EL 23671. The area was considered prospective for skarn-hosted base metal and tin-tungsten deposits. Additionally the Reynolds Range, Lander River valley and Anmatjira Range were considered prospective for hydrothermal skarn uranium deposits.

Otter Exploration commissioned an airborne radiometric survey and identified eighteen radiometric anomalies in the area worth further investigation. Ground truthing of radiometric anomalies was carried out using a scintillometer along traverses. Bore waters in the licence area were analysed for uranium.

Several uranium, thorium and rare earth element (REE) occurrences were located. Otter's exploration activities identified what is now referred to as the Mount Finness REE prospect. The Mount Finness REE occurrence is described by Kojan (1980) as a localised monazite-rich pod about 1-metre in size with activity noted as off-scale (>20,000 CPS).

Further exploration of the Mount Finness REE prospect is unlikely as it lies within an exclusion zone.

BHP Exploration (1981-1983) ELs 2942, 3075, 3084 and 3088.

The tenements were taken up principally in a diamond search, with base metals a secondary consideration. BHP acquired 75 x 20 kg stream sediment samples from favourable trap sites at an average density of about 23 km². BHP withdrew in the light of negative results however all data was not reported in their final report. A coincident stream sediment silt sample was also collected. Two stream sediment samples were anomalous for cerium (Ce) and lanthanum (La): they drained an area near Mt Finness in the north-west corner of EL 23671, and which hosts the Mt Finness REE (Th-U) prospect. This work suggests that stream geochemistry may be useful in a search for further REE deposits in Arafura's tenements. However caution is warranted given the presence of monazite in the surrounding metamorphic rocks.

J Weir (1982-1983) EL3506.

Via a scintillometer search, Weir located an allanite-bearing pegmatite vein near Mt Boothby, which is 5 km north-east of Nolans Bore and within the adjacent EL 23671. The pegmatite is of limited extent and assayed: Ce 4250 ppm, La 3100 ppm and Yttrium (Y) 70 ppm. The presence of REE-enriched pegmatite is noteworthy and encouraging given the presence of calc-silicate country rocks in the Reynolds Range Group.

BHP Exploration (1983-1984) EL4188.

The western half of that tenement lay within the eastern margin of EL 23671. BHP reviewed the BMR 1:250 000 total magnetic intensity contour map for the Napperby Sheet and identified a "bullseye" shaped anomaly of nearly 1 000 nT, the source of which was considered to be a possible carbonatite. It lies just north-east of EL 23671, with a possible repetition inside Arafura's licence. Without any outcrop to explain it, the area was applied for and BHP carried out an aeromagnetic survey and follow-up ground magnetometer and gravity traverses. These surveys confirmed the size and location of the anomaly and identified a five to six milligal gravity anomaly. The geophysical anomalies were drill tested by hole OG-1 (total depth 256 metres) which intersected non-magnetic "basement" rocks at 94 metres, and weakly magnetic hornblende-quartz-feldspar gneiss at 213 metres. It was considered that these rocks belonged to the Arunta Complex and were the source of the geophysical anomaly. Analysis for base metals proved negative.

Colchis Mining Corporation Pty Ltd (1987-1990) EL5511.

This tenement occupied the western part of EL 23671. Colchis targeted both Au and base metal mineralisation. It undertook a review of previous data, interpreted aerial photography and Landsat imagery, and carried out an intensive mapping programme. Rock chip samples (45) and stream sediment sampling (210 samples, not BLEG Au) did not indicate any anomalism warranting Colchis' further work.

Track Minerals Pty Ltd (1988-1989) EL5901.

EL 5901 occupied the southeast part of EL 9672 Dragons Lair and subsequently EL 23671. Track searched for Au in the Aileron Metamorphics, which there consist of a calc-silicate, meta-carbonate and meta-arenite suite. It focused on areas proximal to granitic intrusions, thrust faulting and shearing. Stream sediment sampling with BLEG Au and base metal analysis, geological traverses and rock-chip geochemistry did not provide any encouragement. However, Drummond considers that its 27 sample stream geochemical programme could not be considered an adequate test of the ±300 sq km that Track endeavoured to evaluate.

Poseidon Gold Ltd and Exodus Minerals N L (1993 - 1999) EL7343

The original large tenement was centred on the Lander River valley, and almost all of it lay well north to northwest of EL 23571 Reynolds Range near the Harverson Pass. The target was structurally controlled Au and base metal mineralisation. Strong programmes were undertaken, including airborne geophysical surveys, stream and lag sampling, RAB drilling and geomorphological interpretation, and successive statutory partial relinquishments of the tenement were made. Although Posgold considered that no anomalous geochemical systems had been defined, Exodus states that "... the relinquished portion ... is not unprospective...": this comment, presumably, being applicable to the Harverson Pass area.

Tidegate Pty Ltd (1993-1994) EL8117.

Aileron roadhouse lies in the north-west part of this tenement, which covered the eastern part of EL9672 Dragons Lair and the southerly lobe of EL 23671 Aileron. Work consisted of taking BLEG Au samples, soil samples and rock chip samples near a known nickel (Ni)-chrome (Cr) prospect, and the amphibolite at Harry's Yard (both of these lie to the south-east of Arafura's tenements). Exploration target was Au in greenstones but results were discouraging.

Aberfoyle Resources Ltd (1995-1998) EL9146.

This tenement covered the south-eastern portion of EL 23671 Aileron. Aberfoyle considered that the poorly outcropping sequences in the tenement could host Granites-Tanami style Au mineralisation, and that the latter may be associated with magnetic anomalies. An early RAB drill programme (6 holes, 299 metres) established that Cainozoic cover thickness was not prohibitive (between 12 and 38 metres). A detailed aeromagnetic survey was flown and several magnetic anomalies located which Aberfoyle considered worthy of follow-up. Apparently this was not undertaken, and the tenement was relinquished.

PNC Exploration (Australia) Pty Ltd (1994-1996) EL8411.

PNC acquired EL8411 so that it could seek uranium. It covered Arafura's relinquished tenement EL9672 Dragons Lair and extended north-westerly well beyond it along the Reynolds Range, now EL 23571 Reynolds Range and EL 23671 Aileron. In its exploration it discovered the Nolans Bore apatite-REE deposit. Via regional reconnaissance, airborne radiometrics and ground traversing, PNC found numerous prospects, many of which it considered were of Mary Kathleen or East Alligator style. However, they were apparently too small to justify more detailed work. Despite the discovery of Nolans Bore, PNC relinquished the tenement.

NORTHERN TERRITORY GEOLOGICAL SURVEY

The Napperby-Hermannsberg 1:250,000 map sheet areas were surveyed in late 1997 by the Northern Territory Geological Survey (NTGS). Flight line spacings were 400 metres on a north-south line orientation. Readings were taken at 70 metre intervals for radiometrics and 7 metre intervals for magnetics. The survey was navigated using Differential GPS at an average terrain clearance of 60 metres. Data was collected by the World Geoscience Corporation.

EXPLORATION BY ARAFURA RESOURCES FROM 1996 TO 2003 (after Drummond 2003)

Homestake Gold of Australia Pty. Ltd. (1996 – 1998) EL 9672.

EL 9672 Dragons Lair was farmed out to Homestake Gold of Australia Ltd. Homestake perceived potential for Granites-Tanami style mineralisation associated with a major fluid pathway, the Trans-Tanami structural zone. It was encouraged by the small Au and Cu prospects in the tenement, and by the proximity to the Sabre Province (Cu-Ag & Cu-Pb-Zn).

The ±1500 sq km of the tenement were covered by 235 BLEG Au samples. Results were disappointing, with only 15 samples attaining 0.1 ppb or better. Six of those were clustered near the Harverson Pass, with a maximum value of 34.6 ppb and three others attaining 0.3 ppb or better. The anomalous area was followed up with detailed stream BLEG sampling (114 samples) and 10 chip samples of rocks with encouraging appearance. Essentially all BLEG samples recorded results below the level of detection. Homestake then withdrew, apparently unaware of PNC's Nolans Bore discovery. Drummond (2003) considered a minus 4mm fraction BLEG program, and a follow-up minus 2mm fraction BLEG program was ineffective, accounting for poor duplication in results.

Ironstone occurrences were investigated in the Harverson Pass and upper Woodforde River areas by Lindsay-Park (1998) and Goulevitch (1999). Lindsay-Park (1998) described haematitic beds in a zone 50-100 metres wide at a stratigraphic change from arenitic units, (now mica schist), to coarser arenitic units, (now quartzite) in the Harverson Pass area. The highest assay result for iron was 21.4%. Goulevitch (1999) investigated goethitic ironstone occurrences in the upper Woodforde River area that were surficial developments in a major marble/limestone unit of the Woodforde River Beds, (upper Reynolds Range Group). Assay of the grab samples yielded 46.10% Fe, and slightly elevated base metals.

INVESTIGATIONS BY ARAFURA RESOURCES ON EL 27290

Arafura Resources' activities and investigations of EL 27290 are noted below.

Arafura acquired HyMap hyperspectral survey data and processed imagery, covering the majority of the Aileron-Reynolds project area, which includes EL 27290, in 2008 prior to the grant of EL 27290 (Hussey and Hornibrook 2008). The acquisition of this hyperspectral survey data was seen as a method of refining reconnaissance exploration activities by focussing on specific mineral signatures. EL 27290 occupied a small gap on the southern side of SEL 23671 that was flown across during the 2008 regional hyperspectral survey.

Hussey and Hornibrook (2008) processed the hyperspectral data using standard library signatures for various minerals and identified 10 meaningful end-members within the SWIR. It was decided that ground-truthing of these mineral signatures and refinement to the processed imagery may be required and after reconnaissance sampling.

Targets were generated from the standard hyperspectral image dataset provided by Hussey and Hornibrook (2008) and regional radiometric survey data. The relevant hyperspectral mineral end-members included topaz, kaolinite, chlorite, epidote, carbonate, amphibole (tremolite-actinolite), muscovite, paragonite and apatite/epidote mineral end-members. Combinations of these mineral signatures can form in phyllic, argillic, advanced argillic and propylitic alteration assemblages. In addition and the principal reason for the survey, some of these mineral end-member signatures may also be useful in locating additional Nolans Bore type-mineralisation. Hyperspectral mineral maps for haematite, goethite and dolomite were also used to assist geological knowledge.

In May-September 2009 during the application phase and prior to the granting of EL 27290, Arafura conducted systematic desktop studies to identify radiometric and hyperspectral mineral targets in the Aileron-Reynolds project area using Arafura's data in conjunction with other relevant GIS layers.

Additional desktop assessments and hyperspectral modelling/reprocessing were also conducted in early 2010 to generate new targets for the 2010 field season (Hussey 2010).

As per Arafura's Exploration Agreement, Arafura submitted a detailed work plan to the CLC for clearance to undertake these reconnaissance activities from late 2009 onwards. Arafura Resources was granted Sacred Site Clearance Certificate No. 2009-075 on 25 September 2009. A number of conditions and restricted areas were identified in the region and hence some of the proposed regional reconnaissance sampling was aborted. No restrictions were applied to EL 27290.

Arafura Resources conducted 4WD vehicle-assisted reconnaissance exploration across the Reynolds Range project area in October-November 2009. Preliminary reconnaissance was also conducted on EL 27290 at that time with a total of 6 rock chip samples collected for assay. Samples were subsequently assayed in late December 2009 after the granting of EL 27290.

In addition to these 6 sites, two additional REE hyperspectral target sites (ARA0611 and ARA0612) were visited in early 2010. These targets were generated during reprocessing of the hyperspectral data in 2010. Both sites were found to be a mixture of mulga and colluvium with no radiometric anomolism; suggesting that the hyperspectral REE target signature was a false positive due to low signal response and instrumental noise.

RESULTS/DISCUSSION

The geochemical results of all rock chip assays are compared to the average Upper Continental Crust (UCC) values of Rudnick and Gao (2005). Assay results for each element of interest were sorted and filtered such that the normalised values of about 5-10x UCC, 10-100x UCC and >100x UCC were colour-coded green, yellow and red, respectively (see Appendix 1). Under this scheme, it is suggested that values of 10-100x UCC are “possibly anomalous” while those >100x UCC are “definitely anomalous”. Values of 5-10x UCC may also be of interest. The systematic use of this methodology allows assays of potential geological interest to be quickly determined in a robust scientific fashion by highlighting potential rock types and areas that show significantly elevated background values with respect to crustal averages.

No “definitely anomalous” values were found during exploration of EL27290. Hyperspectral target site ARA0175 contains “possibly anomalous” Bi, P, Se, and Sn values. Of these P and Sn would appear to be the most promising as they suggest the possibility of hydrothermal veins or skarn development in the area. Furthermore it should also be noted that both Y and Ta are also slightly elevated and may be of interest in this sample, consistent with the “possibly anomalous” geochemical signature noted above.

All of the assayed samples have low REE contents. Despite the low REE contents of the assayed samples, the geochemical associations identified in ARA0175 suggest there may be potential for Sn-P-Y-HREE (*ie* cassiterite-xenotime) mineralisation (probably as veins/skarns) in the area. This observation is consistent with the Sn-W-Ta prospects in the region and the abundance of fractionated granites and calcsilicate rocks and warrants further systematic investigations.

Most of EL 27290 is covered by recent colluvium and the potential for Sn-P-Y-HREE mineralisation should not be dismissed. A more systematic assessment of the area is warranted.

Nolan bore-type mineralisation was not been identified on EL 27290 however it should be pointed out that the area is largely covered and exploration activity on this tenement has been limited. A detailed airborne geophysical survey of the area is recommended and considered essential for a thorough evaluation of the mineral potential of the tenement.

REFERENCES/SOURCES OF INFORMATION

Ahmad M and Scrimgeour IR, 2004. Geological map of the Northern Territory, 1:2 500 000 geological map series. *Northern Territory Geological Survey*.

Andrew Drummond and Associates, Independent Consulting Geologists Report for Arafura Resources NL.

Barnes SJ, Anderson JAC Smith TR and Bagas L, 2008. The Mordor Alkaline Igneous Complex, central Australia: PGE-enriched disseminated sulphide layers in cumulates from a lamprophyric magma. *Mineralium Deposita*, **43**, 641-662.

Buick IS and Cartwright I, 1994. The significance of early scapolite in greenschist facies marbles from the Reynolds Range Group, central Australia. *Journal of Geological Society of London*, **151**, pp. 803-812.

Buick IS, Cartwright I & Harley SL, 1998. The retrograde P-T-t path for low-pressure granulites from the Reynolds Range, central Australia; petrological constraints and implications for low-P/high-T metamorphism. *Journal of Metamorphic Geology*, **16**, 511-529.

Cartwright I, Buick IS, Foster DA and Lambert DD, 1999. Alice Springs age shear zones from the southeastern Reynolds Range, central Australia. *Australian Journal of Earth Sciences*, **46**, 355-363.

Buick IS, Miller JA, Williams IS and Cartwright I, 2001. Ordovician high-grade metamorphism of a newly recognised late Neoproterozoic terrane in the northern Harts Range, central Australia. *Journal of Metamorphic Geology* **19**, 373-394.

Buick IS, Hand M, Williams IS, Mawby J, Miller JA and Nicoll RS, 2005. Detrital zircon provenance constraints on the evolution of the Harts Range Metamorphic Complex (central Australia): links to the Centralian Superbasin. *Journal of the Geological Society, London* **162**, 777-787.

Claoué-Long J, 2003. Event chronology in the Arunta Region. In Munson TJ and Scrimgeour I (Editors), Annual Geoscience Exploration Seminar (AGES) 2003. Record of Abstracts. *Northern Territory Geological Survey, Record 2003-001*.

Claoué-Long J, Fraser G, Huston D, Neumann N and Worden K, 2005. Towards a correlation of the earliest Proterozoic evolution in central Australia. In Munson TJ (Editor), Annual Geoscience Exploration Seminar (AGES) 2003. Record of Abstracts. *Northern Territory Geological Survey, Record 2003-001*.

Claoué-Long JC and Hoatson DM, 2005. Proterozoic mafic-ultramafic intrusions in the Arunta Region, central Australia. Part 2: Event chronology and regional correlations. *Precambrian Research* **142**, 134-158.

Claoué-Long J, Edgoose C and Worden K, 2008a. A correlation of Aileron Province stratigraphy in central Australia. *Precambrian Research* **166**, 230-245.

Claoué-Long J, Maidment D, Hussey K and Huston D, 2008b. The duration of the Strangways Event in central Australia: Evidence for prolonged deep crustal processes. *Precambrian Research* **166**, 246-262.

Close D, Scrimgeour I, Duffett M, Worden K and Goscombe B, 2005. East Arunta project – preliminary results and future directions. In Munson TJ (Editor), Annual Geological Exploration Seminar (AGES) 2005, Record of Abstracts. *Northern Territory Geological Survey Record 2005-001*.

Collins WJ, and Teyssier C, 1989. Crustal scale ductile fault systems in the Arunta Inlier, central Australia. *Tectonophysics*, **158**, pp. 49-66.

Collins WJ and Williams IS, 1995. SHRIMP ionprobe dating of short-lived Proterozoic tectonic cycles in the northern Arunta Inlier, central Australia. *Precambrian Research*, **71**, pp. 69-90.

Collins WJ and Shaw RD, 1995. Geochronological constraints on orogenic events in the Arunta Inlier: a review. *Precambrian Research* **71**, 315-346.

Collins WJ and Sawyer E, 1996. Pervasive granitoid magma transfer through the lower-middle crust during non-coaxial compressional deformation. *Journal of Metamorphic Geology*, **14**, pp. 565-579.

Connors G, 2004. NT Parks and Conservation Masterplan – Burt Plain Bioregion Conservation value and Environmental Resources map. *Northern Territory Department of Natural Resources, Environment and the Arts*.

Dirks PHGM and Wilson CJL, 1990. The geological evolution of the Reynolds Range, Central Australia: Evidence for three distinct structural/metamorphic cycles. *Journal of Structural Geology*, **12**, pp. 651-665.

Dirks PHGM, Hand M and Powell R, 1991. The P-T-deformation path for a mid-Proterozoic, low-pressure terrain: the Reynolds Range, central Australia. *Journal of Metamorphic Geology*, **9**, pp. 641-661.

Donnellan N and Johnstone A, 2003. Expanding the Tennant Region: mapped and interpreted geology of the Mount Peake and Lander River 1:250 000 sheets. In TJ Munson and I Scrimgeour (Editors), Annual Geological Exploration Seminar (AGES) 2003, Record of Abstracts. *Northern Territory Geological Survey Record 2003-0001*.

Fabray J, 2005. The Uranium Potential of the Reynolds Range & Yalyirimbi Range Regions (EL 23571, EL 23671 and ELA 24548), Aileron Project. *Exploremin Pty Ltd unpublished report EPL-05/173*.

Ferenczi PA, 2001. Iron ore, manganese and bauxite deposits of the northern Territory. *Northern Territory Geological Survey, Report 13*.

Fraser G, 2004. Defining the “footprint” of tectonothermal events in the North Australian Craton: recent ⁴⁰Ar/³⁹Ar results from the Davenport Ranges and Barrow Creek Regions. In TJ Munson and I Scrimgeour (Editors), Annual Geological Exploration Seminar (AGES) 2004, Record of Abstracts. *Northern Territory Geological Survey Record 2004-001*.

Freeman, M. J., Shergold, J. H., Morris, D. G., and Walter, M. R., 1990. Late Proterozoic and Palaeozoic Basins of Central and Northern Australia. *Geology of the Mineral Deposits of Australia and Papua New Guinea* (Ed. F.E. Hughes), *AusIMM Monograph 14*, 1125-1134.

Goscombe B, 1991. Intense non-coaxial shear and the development of mega-scale sheath folds in the Arunta Block, central Australia. *Journal of Structural Geology*, **13**, pp. 299-318.

Goulevitch, J, 1999. Annual Report for the period 25/11/98-24/11/99 Year 3. EL9672 Dragon’s Lair, Aileron, Northern Territory. *Exploremin Pty. Ltd., unpublished report EPL-99/103*.

Goulevitch J, 2005. Annual Report SEL 23671 Y/E 07/12/04, RC drilling results at Nolans Bore REE/P deposit, Aileron, Northern Territory. *Exploremin Pty Ltd unpublished report EPL 04/168*.

Goulevitch J, 2006. 2005 drilling results and resource estimate Nolans Bore REE/P/U deposit, Aileron, Northern Territory. *Exploremin Pty Ltd unpublished report EPL 05/177*.

Goulevitch J, 2008. Estimate of identified mineral resources as at 7 November 2008, Nolans Bore REE/P/U deposit, Aileron, Northern Territory. *Exploremin Pty Ltd unpublished report EPL 08/184*.

Hallenstein R and Goulevitch J, 2009. Annual Report SEL 23671 Y/E 07/12/08 Aileron, Northern Territory. *Arafura Resources unpublished report ARU-09/002*.

Hallenstein R and Goulevitch J, 2010. Annual Report SEL 23671 Y/E 07/12/09 Aileron, Northern Territory. *Arafura Resources unpublished report ARU-10/001*.

Hand M and Dirks PHGM, 1992. The influence of deformation on the formation of axial planar leucosomes and the segregation of small melt bodies within the migmatitic Napperby Gneiss, Central Australia. *Journal of Structural Geology*, **14**, 591-604.

Hand M, Fanning CM and Sandiford M, 1995. Low-P High metamorphism and the role of high-heat producing granites in the northern Arunta Inlier. *Australian Geological Society, Abstracts*, **40**, 60-61.

Hand M, Mawby J and Miller J, 1999a. U-Pb ages from the Harts Range, central Australia; evidence for early Ordovician extension and constraints on Carboniferous metamorphism. *Journal of the Geological Society, London* **156**, 715-730.

Hand M, Mawby J, Miller JA, Ballèvre M, Hensen B, Möller A and Buick IS, 1999b. The tectonothermal evolution of the Harts and Strangways Range Region. Field Guide 4. Geological Society of Australia, Specialist Group in Geochemistry, Mineralogy and Petrology.

Hand M and Buick IS, 2001. Tectonic evolution of the Reynolds-Anmatjira Ranges: a case study in terrain reworking from the Arunta Inlier, central Australia. In: JA Miller, RE Holdsworth IS Buick and M Hand (Editors). *Continental Reactivation and Reworking*. Geological Society, London, Special Publications **184**, 237-260.

Hoatson DM, Sun Shensu and Claoué-Long JC, 2005. Proterozoic mafic-ultramafic intrusions in the Arunta Region, central Australia. Part 1: Geological setting and mineral potential. *Precambrian Research* **142**, 93-133.

Hussey KJ, Huston DL and Frater M, 2004. Metallogeny in the eastern Arunta Region and the potential of its Palaeoproterozoic rocks. In TJ Munson and I Scrimgeour (Editors), Annual Geological Exploration Seminar (AGES) 2004, Record of Abstracts. *Northern Territory Geological Survey Record* 2004-001.

Hussey KJ, Huston DL and Claoué-Long JC, 2005. Geology and origin of some Cu-Pb-Zn (-Au-Ag) deposits in the Strangways Metamorphic Complex, Arunta Region, Northern Territory. *Northern Territory Geological Survey Report* 17.

Hussey KJ, 2009. Annual Report EL 23571 Y/E 07/12/08 Aileron, Northern Territory. *Arafura Resources unpublished report ARU-09/003*.

Hussey KJ, 2010a. Annual Report EL 23571 Y/E 07/12/08 Aileron, Northern Territory. *Arafura Resources unpublished report ARU-10/002*.

Hussey KJ, 2010b. Report on 2009 reconnaissance program in the Aileron-Reynolds Project area, Northern Territory. *Arafura Resources unpublished report ARU-10/003*.

Hussey MC and Hornibrook MJ, 2008. HyMap Survey and processing report, Reynolds Range, NT. *HyVista Corporation report*.

Kojan CJ, 1979. Annual report Exploration Licence 1444, NT. Otter Exploration NL. *NT Department of Resources Company Report* CR1979-0021.

Kojan CJ, 1980. Exploration Licence 1444, NT. Final report. Otter Exploration NL. *NT Department of Resources Company Report* CR1980-0056.

Kojan CJ and Fortowski D, 1980. 1979 Report on ELs 1581, 1582, 1583, 1584, 1585, 1444, 1445,

1450, 1451, 1702 and 2200 and ELA 2426. Otter Exploration NL. *NT Department of Resources Company Report CR1980-0252.*

Mahar EM, Baker JM, Powell R, Holland TJB and Howell N, 1997. The effect of Mn on mineral stability in metapelites. *Journal of Metamorphic Geology*, **15**, 223-238.

Maidment DW, Williams IS and Hand M, 2004. The Harts Range Metamorphic Complex – a Neoproterozoic to Cambrian rift sequence in the eastern Arunta Region. In TJ Munson and I Scrimgeour (Editors), Annual Geoscience Exploration Seminar (AGES) 2004. Record of Abstracts. *Northern Territory Geological Survey Record 2004-001.*

Mawby J, Hand M, Foden J and Kinny P, 1998. Ordovician granulites in the eastern Arunta Inlier: a new twist in the Palaeozoic history of central Australia. *Geological Society of Australia, Abstracts* 49, 296.

Mawby J, Hand M and Foden J, 1999. Sm-Nd evidence for Ordovician granulite facies metamorphism in an intraplate setting in the Arunta Inlier, central Australia. *Journal of Metamorphic Geology* 17, 653-668.

Miller JA, Buick IS, Williams IS and Cartwright I, 1998. Re-evaluating the metamorphic and tectonic history of the eastern Arunta Block, central Australia. *Geological Society of Australia, Abstracts* 49, 316.

McGivray T, 2006. Helicopter-borne reconnaissance October, 2005 EL 23571 Reynolds Range Northern Territory. Annual report EL 23571 for year ending 07/12/05, Reynolds Range project, Northern Territory. *Arafura Resources NL, Report ARU 2006-004.*

Naldrett AJ, 2004. *Magmatic Sulfide Deposits: Geology, Geochemistry and Exploration.* Springer.

Rubatto D, Hermann J and Buick IS, 2006. Temperature and bulk composition control on the growth of monazite and zircon during Low-pressure anatexis (Mount Stafford, Central Australia). *Journal of Petrology*, **47**, 1973-1996.

Rudnick RL and Gao S, 2005. Composition of the Continental Crust, pp 1-64. In: *The Crust* (RL Rudnick, Editor) Volume 3 Treatise on Geochemistry. (HD Holland and KK Turekian, Editors), Elsevier, Oxford.

Scrimgeour I, 2003. Developing a revised framework for the Arunta Region. In TJ Munson and I Scrimgeour (Editors), Annual Geological Exploration Seminar (AGES) 2003, Record of Abstracts. *Northern Territory Geological Survey Record 2003-001.*

Scrimgeour I and Raith JG, 2001. High-grade reworking of Proterozoic granulites during Ordovician intraplate transpression, eastern Arunta Inlier, central Australia. In. JA Miller, RE Holdsworth IS Buick and M Hand (Editors). *Continental Reactivation and Reworking.* Geological Society, London, Special Publications 184, 261-287.

Senior, B. R., Truswell, E. M., Iduurm, M., Shaw, R. D., and Warren, R. G., 1995. Cainozoic sedimentary basins in the eastern Arunta Block, Alice Springs Region, central Australia. *Australian Geological Survey Organisation - Journal of Australian Geology & Geophysics*, **15(4)**, pp. 421-444.

Shaw RD, Stewart AJ and Black LP, 1984. The Arunta Inlier: a complex ensaillc mobile belt in central Australia. Part 2: tectonic history. *Australian Journal of Earth Sciences*, 31, 457-484.

Stewart, A. J., Offe, L. A., Gliksn, A. J., Warren, R. G., and Black, L. P., 1980. *Geology of the northern Arunta Block, Northern Territory.* Australian Bureau of Mineral Resources, Geology and Geophysics Record, **1980/63.**

Stewart AJ and Pillinger DM, 1981. Reynolds Range Region, Northern Territory. *BMR 1:100,000 Geological Map Series Commentary.* Bureau of Mineral Resources, Geology and Geophysics,

Canberra, Australia.

Stewart AJ, Shaw RD and Black LP, 1984. The Arunta Inlier: a complex ensaillc mobile belt in central Australia. Part 1: stratigraphy, correlations and origin. *Australian Journal of Earth Sciences*, 31, 445-455.

Stewart, A. J., 1991. NAPPERBY – Sheet SF/53-9. *BMR 1:250,000 Geological Map Series and Commentary*. Bureau of Mineral Resources, Geology and Geophysics, Canberra, Australia

Sun Shensu, Warren RG and Shaw RD, 1995. Nd isotope study of granites from the Arunta Inlier, central Australia: constraints on geological models and limitation of the method. *Precambrian Research* 71, 301-314.

Thevisson, J, 1995. Napperby Annual Report EL8411 1995 field season. PNC Exploration (Australia) Pty Ltd unpublished report. *NT Department of Resources Company Report CR1996/0187*.

Vicary, MJ, 1979. Annual Report 1978/79 on geological exploration EL 1294, Napperby, Northern Territory. CSR Minerals and Chemical Division. *NT Department of Resources Company Report CR1979-0198*.

Vry JK and Cartwright I, 1998. Stable isotopic evidence for fluid infiltration during contact metamorphism in a multiply-metamorphosed terrane: the Reynolds Range, Arunta Block, central Australia. *Journal of Metamorphic Geology*, 16, 749-765.

Vry JK, Compston W and Cartwright I, 1996. SHRIMP II dating of zircons and monazites: reassessing the timing of high-grade metamorphism and fluid flow in the Reynolds Range, northern Arunta Block, Australia. *Journal of Metamorphic Geology*, 14, 335-350.

Wells, R. G., 1972. Annual Report, 1972. *Tanganyika Holdings Pty. Ltd. Unpublished Company Report: CR1972-0063*.

Wells AT and Moss FJ, 1983. The Ngalia Basin, Northern Territory: stratigraphy and structure. *Bur. Min. Res. Geol. & Geophys., Bulletin* 212, 4-7.

Worden KE, Carson CJ, Close DF, Donnellan N and Scrimgeour IR, 2008. Summary of results. Joint NTGS-GA geochronology: Tanami Region, Arunta Region, Pine Creek Orogen and Hallc Creek Orogen correlatives, January 2005- March 2007. *Northern Territory Geological Survey, Record* 2008-003.

Williams IS, Buick IS and Cartwright I, 1996. An extended episode of early Mesoproterozoic metamorphic fluid flow in the Reynolds Range, central Australia. *Journal of Metamorphic Geology*, 14, 29-48.

Xu G, Will TM and Powell R, 1994. A calculated petrogenetic grid for the system K₂O-FeO-MgO-Al₂O₃-SiO₂-H₂O with particular reference to contact-metamorphosed pelites. *Journal of Metamorphic Geology*, 12, 99-119.

Zhao Jianxin and Bennett VC, 1995. SHRIMP U-Pb zircon geochronology of granites in the Arunta Inlier, central Australia: implications for Proterozoic crustal evolution. *Precambrian Research* 71, 17-43.

Zhao Jianxin and McCulloch MT, 1995. Geochemical and Nd isotopic systematics of granites from the Arunta Inlier, central Australia: implications for Proterozoic crustal evolution. *Precambrian Research* 71, 265-299.