

ANNUAL REPORT

ON EXPLORATION WORK CARRIED OUT ON

E.L. 105 BYNOE BAY

NORTHERN TERRITORY

APPENDIX ONE - Assay Results and Location Plans

APPENDIX TWO - Airborne Maps and Report

## CONTENTS

1. INTRODUCTION
2. WORK DONE
3. GEOLOGY
4. GEOPHYSICS
5. GEOCHEMISTRY
6. EXPENDITURE
7. CONCLUSION

## INTRODUCTION

E.L. 105 covering a total area of 102 square miles in the Bynoe Bay area was granted to Mr. P.E. Cogar on 7/5/72 for a period of twelve months. This area was covered by Authority to Prospect 2751 prior to conversion to its present E.L. status.

Field work in this area is hampered by its excessive wet nature, heavy scrub, soakages and poor access. In addition, the Cretaceous cover generally up to 50' thick covers most of the Lower Proterozoic and Archaean rock types.

Expenditure committment for this year was \$5,000.00.

Application for renewal of this E.L. was lodged 3/5/73.

## WORK DONE

The work carried out on this licence has consisted primarily of an airborne gamma ray spectrometer and total magnetic intensity survey. The contract for the airborne survey was let to Aero Service (Australia) Pty. Ltd. Data interpretation was done by Layton and Associates Pty. Ltd. Based on the results, a number of field trips were made by the company's consultants to the more accessable anomalous areas along the northern boundary of the E.L. Although anomalies designated in the report as A3 and A4 have not been checked out to date, the large total count anomalies that have been evaluated are caused by anomalous thorium content.

## GEOLOGY

A number of geological traverses were made along the more accessable northern boundary of the E.L. which covers from the nearby Raft Point to Pioneer Beach. The remainder of the E.L. is extremely difficult to evaluate as the shoreline of Bynoe Bay is lined by mangrove swamps.

Near Raft Point an Archaean sequence of gneiss and schist intruded by the Litchfield Intrusive Complex is overlain by relatively flat lying, up to 50' thick, Cretaceous quartz sandstone. Along the flanks of the Litchfield Intrusive outcrops, the Cretaceous consists of near residual angular to sub-angular quartz pebble conglomerate.

A total count anomaly designated A5 just north of Keswick Point along the northern boundary was investigated (see airborne geophysical survey interpretation plan). A number of shallow costeans were bulldozed over the area in addition to those sunk by Thiess Brothers and Australus Mining. These costeans exposed a weathered quartz - feldspar porphyry with pegmatitic phases. In one costean (see plan) a total count of 1500 to 2500 cps was registered over a narrow 2" - 3" band to 2' in depth within a laterized zone (see sample F7677). No secondary or primary

mineralization has been found within the anomalous area.

In addition, a number of auger holes were drilled to a depth of 9' with samples taken at 3' and 9'.

The radioactivity probably originates either from one of the thorium - cesium silicates or from the phosphate mineral monazite.

At Pioneer Beach, badly weathered granodiorite (?) crops out, and is regarded to be part of the Litchfield Intrusive Complex.

#### GEOPHYSICS

An airborne gamma ray spectrometer and total magnetic survey was flown and compiled by Aero Service (Australia) Pty. Ltd. during late November. Layton Geophysical Consultants Pty. Ltd. interpreted the data and their report plus plans are attached for your information. Twelve total count anomalies have been delineated and designated by prefix A. Anomalies A3 and A4 have little or no associated thorium response. Follow-up work is warranted in particular to the latter.

Ground scintillometer survey was conducted over anomaly designated as A5. Background was established as being 50 cps. This survey outlined a 500 metre wide N-S trending anomaly of greater than 100 cps. This zone appears to outline a weathered near surface quartz - feldspar granite with pegmatitic phases. (See sketch plan).

#### GEOCHEMISTRY

Seven grab samples were taken from existing and recent costeans sunk by Kewanee Australia Pty. Ltd., from the more anomalous total count zones. (See sketch plan for location).

Sample No.	Th.	$U_3O_8$
F 7676	184	330
7677	3770	120
7678	199	4
7679	783	26
7680	281	10
7681	148	4
7682	193	-

Also attached are results from 21 angle holes to a depth of 9' and sampled at 3' and 9'.

The results confirm the airborne survey related to thorium bearing minerals.

At Pioneer Beach, a number of grab samples were taken over anomalous total count areas as outlined by scintillometer survey. Results are attached.

Assay results to date confirm airborne survey data that with the possible exception of anomalous areas designated A3 and A4, radio activity is largely related to thorium bearing rock types, probably later pegmatites within the Litchfield Intrusive Complex.

#### EXPENDITURE

The expenditure on the area which includes time spent by Consultants on this licence, data and airborne interpretation, and related office overheads amounts to \$6,000.00.

#### CONCLUSION

The results of the airborne survey are encouraging with respect to the number of anomalies that have been outlined in particular A3 and A4. However, it is quite possible that the large total count anomalies are largely caused by Thorium bearing minerals.

Signed on behalf of  
Kewanee Australia Pty. Ltd.

BY:

  
J.B. Felderhof  
Consulting Geologist.

Bynoe way.

LABOR... 19...

Th

386

LAMSON PARAGON (M-1) LIMITED - FORMALINER PARAFLO

				Th
1				
2				
3	7683F	26644	Hole 1	3' 0037
4				
5	7684F	26645	Hole 1	9' 0083
6				
7	7685F	26646	Hole 2	3' 0020
8				
9	7686F	26647	Hole 2	9' 0006
10				
11	7687F	26648	Hole 3	3' 0033
12				
13	7688F	26649	Hole 3	9' 0101
14				
15	7689F	25550	Hole 4	3' 0019
16				
17	7690F	26651	Hole 4	9' 0009
18				
19	7691F	26652	Hole 5	3' 0037
20				
21	7692F	26653	Hole 5	9' 0031
22				
23	7693F	26654	Hole 6	3' 0025
24				
25	7694F	26655	Hole 6	9' 0044
26				
27	7695F	26656	Hole 7	3' 0086
28				
29	7696F	26657	Hole 7	9' 0046
30				
31	7697F	26658	Hole 8	3' 0042
32				
33	7698F	26659	Hole 8	9' 0106
34				
35	7699F	26660	Hole 9	3' 0094
36				
37	7700F	26661	Hole 9	9' 0083
38				
39	7701F	26662	Hole 10	3' 0048
40				
41	7702F	26663	Hole 10	9' 0052
42				
43	7703F	26664	Hole 11	3' 0026
44				
45	7704F	26665	Hole 11	9' 0024
46				
47	7705F	26666	Hole 12	3' 0038
48				
49	7706F	26667	Hole 12	9' 0128
50				
51	7707F	26668	Hole 13	3' 0073
52				
53	7708F	26669	Hole 13	9' 0067
54				
55	7709F	26670	Hole 14	3' 0127
56				
57				
58				
59				
60				

26

Bynoe Bay.

M. S. N. Laboratories Pty. Ltd.

Th

2864

		Th
1		
2		
3	7683F	26644
4		
5	7684F	26645
6		
7	7685F	26646
8		
9	7686F	26647
10		
11	7687F	26648
12		
13	7688F	26649
14		
15	7689F	26650
16		
17	7690F	26651
18		
19	7691F	26652
20		
21	7692F	26653
22		
23	7693F	26654
24		
25	7694F	26655
26		
27	7695F	26656
28		
29	7696F	26657
30		
31	7697F	26658
32		
33	7698F	26659
34		
35	7699F	26660
36		
37	7700F	26661
38		
39	7701F	26662
40		
41	7702F	26663
42		
43	7703F	26664
44		
45	7704F	26665
46		
47	7705F	26666
48		
49	7706F	26667
50		
51	7707F	26668
52		
53	7708F	26669
54		
55	7709F	26670
56		
57		
58		
59		
60		

LAUSON PARACON (P) LIMITED - FORMLINER PARAFLO

26

duplicate

BYNOE

BAY

AUGER. SAMPLES

				Th		
1					U3 07	
2	7901F	LINE 1 W	NO. 22	DEPTH 3'	0087	0004
4	7902F	1 W	22	9'	0138	0005
6	7903F	1 W	23	3'	0124	0003
8	7904F	1 W	23	9'	0088	0014
10	7905F	1 W	24	3'	0092	0006
12	7906F	1 W	24	9'	0058	0004
14	7907F	1 W	25	3'	0029	0006
16	7908F	1 W	25	9'	0040	0008

8

33 total  
 52  
 85

BYMORE BAY  
 AUGER. SAMPLES  
 U<sub>3</sub>O<sub>8</sub>.

			DEPTH	Th
1	7710F	B. B. LINE 1 W	HOLE 14. 9'	0070
2	7711F	" 1 W	" 15 3'	0150
3	7712F	" 1 W	" 15 9'	0110
4	7713F	" 1 W	" 16. 3'	0136
5	7714F	" 1 W	" 16 9'	0204
6	7715F	" 1 W	" 17 3'	0094
7	7716F	" 1 W	" 17 9'	0652
8	7717F	" 1 W	" 18. 3'	0222
9	7718F	" 1 W	" 18 9'	0078
10	7719F	" 1 W	" 19 3'	0158
11	7720F	" 1 W	" 19 9'	0206
12	7721F	" 1 W	" 20 3'	0219
13	7722F	" 1 W	" 20 9'	0155
14	7723F	" 1 W	" 21 3'	0110
15	7724F	" 1 W	" 21 9'	0033

0002  
 0005  
 0004  
 0010  
 0008  
 0006  
 0012  
 0007  
 0003  
 0006  
 0006  
 0010  
 0007  
 0004  
 0005

Ass

Re

15

BYMORE BAY  
 AUGER. SAMPLES  
 U<sub>3</sub>O<sub>8</sub>.

			DEPTH	Th	
7710F	B. B. LINE 1 W	HOLE	14. 9'	0070	0002
7711F	"	"	15 3'	0150	0005
7712F	"	"	15 9'	0110	0004
7713F	"	"	16. 3'	0136	0010
7714F	"	"	16 9'	0204	0008
7715F	"	"	17 3'	0094	0006
7716F	"	"	17 9'	0652	0012
7717F	"	"	18. 3'	0222	0007
7718F	"	"	18 9'	0078	0003
7719F	"	"	19 3'	0158	0006
7720F	"	"	19 9'	0206	0006
7721F	"	"	20 3'	0219	0010
7722F	"	"	20 9'	0155	0007
7723F	"	"	21 3'	0110	0004
7724F	"	"	21 9'	0033	0005

Ass.

Duplicated.

Re.

			Th	U <sub>3</sub> O <sub>8</sub>	Assay
25938	Soil		0196	0016	
25939	-100 mesh		0166	0010	
25940	+100 mesh		0148	0019	
25941	Sand		0424	0046	
85943	+100 mesh		0420	0066	
25944	-100 mesh	more than	1000	0200	

36.60

• Samples taken from Pioneer Beach, Byron Bay.

Results in p

J. Woodin

6

BYNOE BAY

AUGER SAMPLES

				Th	U <sub>3</sub> O <sub>8</sub>	As
1						
2	7901F	LINÉ 1 W	NOLE 22	DEPTH 3'	0087	
4					0004	
6	7902F	1 W	22	9'	0138	
8					0005	
10	7903F	1 W	23	3'	0124	
12					0003	
14	7904F	1 W	23	9'	0088	
16					0014	
18	7905F	1 W	24	3'	0092	
20					0006	
22	7906F	1 W	24	9'	0058	
24					0004	
26	7907F	1 W	25	3'	0029	
28					0006	
30	7908F	1 W	25	9'	0040	
32					0008	

Duplicated.

Res

		$^{232}\text{Th}$	$\text{U}_3\text{O}_8$
25938	Soil	0196	0016
25939	-100 mesh	0166	0010
25940	+100 mesh	0148	0019
25941	Sand	0424	0046
25943	+100 mesh	0420	0066
25944	-100 mesh	more than	1000 0200

36.60


Results in

• Samples taken from Pioneer Beach, Byron Bay.

J. Woodie

Pioneer Beach.

6




**SYMBOLS**

-  COSTEAN.
-  150
-  100
-  RAFT POINT RD.
-  TRACK.
-  AUGER DRILLED.

KEWANEE AUSTRALIA PTY. LTD.	
AREA ALONG NORTH BOUNDARY OF BYNOE BAY E.L.	
EL. NO. 105.	
Scale: 1" = 100 m.	Project: 19
Drawn: Sally	Prospect: Leases 1 → 7
Date: 26-7-73.	<b>EL. 105.</b>
Drg. No. 19 — 3.	

LAYOUT BY: J. FELDERHOF.



**SYMBOLS**

- COSTEAN
- 150 C.P.S.
- 100 C.P.S.
- RAFT POINT RD.
- TRACK.
- AUGER DRILLED.

KEWANEE AUSTRALIA PTY. LTD.	
AREA ALONG NORTH BOUNDARY OF BYNOE BAY E.L.	
EL. NO. 105.	
Scale: 1" = 100 m.	Project: 19
Drawn: Sally	Prospect: Leases 1 → 7
Date: 26-7-73.	<b>EL. 105.</b>
Drg. No. 19 — 3.	

LAYOUT BY: J. FELDERHOF.



REFERENCE

QUATERNARY	Qa/Qz	Alluvial soil cover - residual soils
LOWER PROTEROZOIC	Pgl	LICHFIELD COMPLEX
	Pda	Basic intrusives?
	Pln	NOLTEMUS FORMATION

LEGEND

Probable subsurface boundary of basic intrusives as suggested by magnetic contour maps

Magnetic trend

Fault location (possible) inferred from magnetics

Geological outcrop boundary inferred from total count map

A1 U 20 cps  
 8 70 cps  
 U Uranium counts above background  
 K Potassium  
 Th Thorium

T1 Total count anomaly trend

AIRBORNE GEOPHYSICAL SURVEY  
 BYNOE BAY  
 6° 18' 30" S 130° 37' 30" E  
 NORTHERN TERRITORY  
 KEVANEE (AUSTRALIA) PTY. LTD.  
 INTERPRETATION PLAN

SCALE 1:25,000

INTERPRETATION BY LAYTON GEOPHYSICAL CONSULTANTS PTY. LTD.  
 FOR AERD SERVICE AUSTRALIA PTY. LTD.

12 37 S


12 40 S

12 30 S

130 30 E

130 40 E

130 50 E



FLIGHT ALTITUDE 300 FEET  
 FLIGHT INTERVAL TRAVERSE 50 NAUT MILES 1/4 LINE 4 MILES  
 CONTOUR INTERVAL TOTAL MAGNETIC INTENSITY 2 GAMMAS  
 CONTOUR INTERVAL TOTAL COUNT 55 C.P.S.  
 PLANIMETRIC SOURCE TOPOGRAPHICAL MAPS  
 SURVEYED AND COMPILED NOV - DEC 1974  
 A.S.C. PROJECT NO.

AIRBORNE GEOPHYSICAL SURVEY  
 BYNOE BAY  
 EL 105  
 NORTHERN TERRITORY  
 KEWANEE (AUSTRALIA) PTY LTD  
**TOTAL MAGNETIC INTENSITY**  
 REGIONAL FIELD REMOVED  
 SCALE 1:25,000



DRAWN AND COMPILED BY GERO SERVICE INDUSTRIES LTD  
 GERRARD, VIC.

1237S


1238S

1239S

130.30E

130.40E

130.50E



FLIGHT ALTITUDE 10000 FT  
 FLIGHT INTERVAL 1000 FT  
 CONTOUR INTERVAL 100 FT  
 MAGNETIC SOURCE  
 SURVEYED AND COMPILED BY  
 DATE PROJECT NO.

AIRBORNE GEOPHYSICAL SURVEY  
 BYNOE BAY  
 EL 105  
 NORTHERN TERRITORY  
 KEWANEE (AUSTRALIA) PTY LTD  
**TOTAL COUNT**  
 SCALE 1:25000

PRINTED AND PUBLISHED BY  
 KEWANEE (AUSTRALIA) PTY LTD  
 100 RIVERSIDE DRIVE  
 SYDNEY NSW 2000  
 AUSTRALIA

