



Cameco Australia Pty Ltd

LIVERPOOL PROJECT

EL 2855

PARTIAL RELINQUISHMENT REPORT

Date: **October 2005**

Report No.: **LP05-03**

Period: **25 July 2000 to 24 July 2005**

Authors: **Llyle Sawyer, Geologist**
 Geoff Beckitt, Geophysicist
 Gavin Otto, Project Geologist

Copies: **Cameco Corporation (1)**
 Cameco Australia Pty Ltd (1)
 DBIRD Mines and Energy (1)
 Northern Land Council (1)

SUMMARY

This report is the final report for exploration work undertaken on relinquished blocks within the Liverpool Project during the period 25 July 2001 to 24 July 2005. The tenement is part of a joint venture between Cameco Australia Pty Ltd (Cameco) and the Ngalangak Aboriginal Corporation

Exploration activities included airborne magnetic and radiometric surveys, as well as a hyperspectral survey covering the exploration licence. Regional sandstone traverse sampling was conducted with samples being analysed for a suite of geochemical elements and PIMA clay determinations.

Radiometric anomaly ground truthing failed to locate any significant anomalies, with most being attributed to element scavenging by lateritisation of the ferruginous units or Cretaceous sediments. Illite and kaolinite were the dominant clay mineral assemblages noted and is consistent with regional clays in the Gumarrirbang and Marlgowa Sandstone Members of the Kombolgie Subgroup.

TABLE OF CONTENTS

SUMMARY	i
List OF FIGURES	ii
List OF TABLES	ii
List OF Appendices	iii
INTRODUCTION	1
Location and Access	1
Tenure	1
Physiography.....	2
Tenement Geology.....	2
Regional Structure and Geological History	2
Exploration Target	3
Previous Exploration.....	3
EXPLORATION PROGRAM.....	3
Airborne Geophysics	4
Airborne Surveys	4
Sandstone Traverse Sampling.....	5
CONCLUSIONS/DISCUSSION.....	6
BIBLIOGRAPHY	7

LIST OF FIGURES

Location Map	1
EL 2855 Relinquished Block Map.....	2
Airborne Magnetism Map – TMI 1 st Vertical Derivative	4
Airborne Radiometrics Map – Uranium Channel	4
Airborne Radiometrics Map – Thorium Channel	4
Airborne Radiometrics Map – Potassium Channel.....	4
Airborne Radiometrics Map – Total Counts.....	4
Airborne Radiometrics Map – U-Th-K as RGB	4
Location of Hyperspectral Surveys.....	5
Hymap Mark I - AMS Clay Intensity Image	5
Sample Location Map – Geology	5
Sample Location Map - Landsat.....	5
Sample PIMA TSA Clay Distribution Map.....	6

LIST OF TABLES

Summary of Exploration Work Completed During Reporting Period	4
Sample Locations.....	5
Sample Descriptions	5
Sample Lithology and Physical Properties	5
Sample Alteration and Structural Measurements	5
Sample G400 Geochemistry	6
Sample TSA Analysis (Identified Mineral Species).....	6

LIST OF APPENDICIES

DeBeers Final Report.....	4
Hymap Mark I Survey Logistics Report.....	5
Cameco Standard Outcrop Sampling and Processing Procedures.....	6

INTRODUCTION

This report describes program activities carried out during the 2001 to 2005 field seasons on behalf of a joint venture between Cameco Australia Pty Ltd (Cameco) and the Ngalangak Aboriginal Corporation. Since the Exploration Licence is located on Aboriginal Reserve, the exploration program was carried out under the terms of consent documentation agreed with the Northern Land Council pursuant to the Aboriginal Land Rights (Northern Territory) Act 1976.

The program as described, represents the five years of exploration within the blocks relinquished on tenement EL 2855 by the Joint Venture.

The only activity for the first year was the flying of a fixed wing airborne survey by UTS Geophysics of Perth. The survey, which was originally planned for October 2000, was postponed and eventually flown in June 2001.

A hyperspectral survey over the entire tenement was flown in October 2000.

During the period of exploration, helicopter assisted sandstone sampling programs were conducted over EL 2855. 72 outcrop samples being taken from within the relinquished area, over the period of three years of ground based activities.

A Joint Venture agreement was established with DeBeers in 2001, with DeBeers discontinuing the agreement in 2003 after discouraging results from their stream sediment sampling program.

Location and Access

Exploration Licence 2855 is located in central western Arnhem Land. The tenement is centred about 70 km southeast of Nabarlek and 100 km east of Jabiru. It is situated approximately 120 km southeast of the King River project.

The project area has extremely limited vehicular access. A four-wheel drive track is indicated as traversing the extreme south east of the tenement and another accesses the Marlgowa Outstation on the Mann River. All work was performed with helicopter support.

Location Map

Tenure

EL 2855 was granted on the 25th July 2000 for an initial period of six years. On granting, the total area covered by the licence is 1255 km² with 213 km² being excluded from exploration.

In July 2003, a total of 149 blocks (498.7 km²) were relinquished from the original 375 blocks, with 226 blocks (756.3 km²) retained.

On 25 July 2005, 125 blocks (418.6 km²) were relinquished as per requirement under Sections 26 and 27 of the Mining Act. 101 blocks (336.7 km²) were retained. The relinquished portion of EL 2855 is shown in the following figure.

EL 2855 Relinquished Block Map

Physiography

The tenement consists predominantly of relatively flat-lying sandstone plateau covered by savannah woodland and scrubland. The plateau is dissected by numerous, mostly north flowing drainages including the Liverpool and Mann Rivers. Gorges and waterfalls have developed in places.

Tenement Geology

Based on the NTGS mapping of the Milingimbi 1:250000 geological series (Carson and others 1999), the tenement is dominated by the Kombolgie Subgroup (formerly Kombolgie Formation) of the Paleoproterozoic Katherine River Group. The latter forms an extensive area of platform cover sediments in western Arnhem Land. The Marlgowa Sandstone, which is the stratigraphically highest unit of the Kombolgie, predominates with lesser exposures of the McKay Sandstone. The former consists of coarse to occasionally pebbly white-grey quartz arenite and was formed in a fluvial depositional environment. The McKay, which lies conformably on the Marlgowa, consists of interbedded fine to medium grained white-grey quartz arenite and red-brown to purple ferruginous sandstone, occasionally pebbly. The depositional environment varied from fluvial to shallow tidal marine.

The Gilruth Volcanic Member is present over a very limited area in the extreme south western corner of the tenement. This unit, represented by ferricrete and saprolite rubble, separates the Marlgowa from the underlying Gumarrinbang Sandstone. Isolated outcrops of Oenpelli dolerite have been mapped within the tenement. Stratigraphically, these sandstone formations are located at the top of the Kombolgie Subgroup. Depth to basement from the top of the Marlgowa Sandstone on the Milingimbi sheet area is estimated to be at least 700 metres, calculated from the measurement of sections through the various sandstone units by NTGS.

Regional Structure and Geological History

The early Proterozoic rocks of the region have been affected by the Top End orogeny (1880 to 1780 Ma), which includes the initial Nimbuwah Event or Barramundi Orogeny at about 1870 Ma. This produced a prograde metamorphic effect with associated tight folding and faulting. The various 'domains' exhibited a variability of deformation and metamorphic grade with the western and eastern margins of the Pine Creek Inlier (Litchfield Province and Nimbuwah domain respectively) exhibiting the most pronounced effects.

Major regional faults, which affect the early Proterozoic, have northwest (Bulman), north-north-west and northerly (Goomadeer) strikes. Another significant set trends to the east and includes both the Ranger and Beatrice faults. The Bulman Fault Zone is the

principle regional feature and is considered to represent a long-lived deep crustal structure, which has exerted a large lateral component in rocks of the Pine Creek Inlier.

A more intense concentration of structures traverse the mid Proterozoic and younger rocks and include northwest, east, northeast and north trends. Both faulting and jointing with displacements ranging from a few metres up to 100 metres, locally heavily dissect the Kombolgie.

In a regional context, the Liverpool project is located adjacent to the Arnhem Shelf at the northern extent of the McArthur Basin. The tectonic environments that existed during deposition of the Katherine River Group varied, ranging from extension and local basin formation with probable fault-controlled sedimentation (Mamadawerre Sandstone), to a basin-wide extensional setting. As noted above the exposed sandstone units illustrate spectacular eroded joint and fault patterns, however the near horizontal to shallow-dipping bedding would imply a tectonically inactive post depositional environment.

The widespread Oenpelli Dolerite intrusive event took place at about 1715 Ma. Localised effects in the sandstone include silicification, the introduction of magnesium-rich to intermediate chlorite and the formation of muscovite-illite.

Exploration Target

The focus of the exploration strategy is the discovery of unconformity-related uranium deposits. The nearby economic deposits at Ranger, Jabiluka, Koongarra and the now depleted Nabarlek Mine serve as models for this strategy. The presence of gold, palladium and platinum in these deposits plus the economic gold-platinum resource at Coronation Hill in the South Alligator Valley, indicates an additional potential for this deposit style.

Previous Exploration

No exploration has occurred on the Liverpool Project prior to granting of EL 2855.

EXPLORATION PROGRAM

Data from the hyperspectral Hymap Mark I survey, flown in 2000, was processed and interpreted early in the current lease year. Analysis of the interpreted data revealed several zones of high clay content. Digital data for the relinquished area of this survey has been submitted on two accompanying DVD's with this report

During the 2001 field season a fixed-wing magnetic, radiometric and DTM (digital terrain model) survey was flown over the entire project area. Digital data for the relinquished area of this survey has been submitted on CD with this report.

Later helicopter assisted regional sandstone sampling program was conducted over the area of the relinquished blocks. This resulted in the collection of 72 for geochemical assay, clay estimation analysis (PIMA).

A Joint Venture agreement was established with DeBeers in 2001. This agreement entitled DeBeers to conduct diamond exploration over the Liverpool Project. 210 samples were collected during 2002, of which 72 samples are within the relinquished portion of the exploration licence. The full DeBeers report is included as an appendix to this relinquishment report.

DeBeers Final Report

Digital data for the surveys and that available for the sample analyses has been submitted on CDs with this report.

The Summary of Expenditures table is the expenditure statistics itemised by work and exploration license.

Summary of Exploration Work Completed During Reporting Period

Airborne Geophysics

During July 2001, Universal Tracking Systems Pty Ltd (UTS) conducted a single detailed airborne magnetic, radiometric and DTM survey totalling 6860 line km. Survey specifications employed for this survey were 60m flight height, 200m lines flown in a west east direction.

The primary aims of the survey were to locate any surface uranium enrichment and to acquire data useful for future geological mapping. No anomaly identification or geology interpretation has been undertaken within the current reporting period.

The gridded results of airborne geophysical surveys flown over relinquished areas are displayed in the following Figures. The areas excluded from exploration (no-go areas) have been greyed-out by request from Traditional Owners through the Northern Land Council.

Airborne Magnetics Map – TMI 1st Vertical Derivative

Airborne Radiometrics Map – Uranium Channel

Airborne Radiometrics Map – Thorium Channel

Airborne Radiometrics Map – Potassium Channel

Airborne Radiometrics Map – Total Counts

Airborne Radiometrics Map – U-Th-K as RGB

Airborne Surveys

In 2000 a 1255km² hyperspectral survey was flown over the Kombolgie Sandstone outcrops within EL2855. The survey was conducted by De Beers using their AMS (Airborne Multi-spectral Scanner) which has recently been renamed Hymap Mk1. The survey was designed to map minerals and identify alteration associated with

unconformity uranium mineralisation. In particular, it was hoped that this system would identify and map variations in kaolinite, illite, dickite, halloysite, iron and magnesium chlorites and silicification, which could be attributed to the above alteration phenomena.

As described in the logistics report, the data has been submitted in ENVI BIL (Band Interleaved by Line) format because ENVI is the usual processing package for hyperspectral data and allows the survey parameters to be stored. This data can be imported into ERMapper if required.

Location of Hyperspectral Surveys

Hymap Mark I Survey Logistics Report

Hymap Mark I - AMS Clay Intensity Image

The data indicates that the dominant clay layers within the Kombolgie are (in order from bottom to top) illite and kaolinite, dickite, and illite. Alteration is suggested by muscovite plus illite in zones along faults, particularly in the west, and at the edges of disturbed blocks. The processing has not identified chlorite alteration.

Sandstone Traverse Sampling

The helicopter assisted regional sampling programs conducted during the period of ground based field activities from 2002 to 2004 were designed as follow-up ground truthing to the airborne surveys over the Kombolgie Subgroup outcrop, ground truthing of the airborne hyperspectral clay interpretation anomalies and also regional based sampling. A total of 72 samples were collected during the period. The following figure shows the areas of coverage. These samples were selected to represent anomaly and regional background signatures for lithological, spectral and geochemical parameters at each location. Geomorphological, geological and radiometric parameters were recorded and a digital photograph at each site was taken. The samples were systematically processed after return to Darwin. Lithological textures, alteration colours (Munsell), grain-size variations and petrophysical parameters (magnetic susceptibility) were routinely recorded.

Sample Locations

Sample Descriptions

Sample Lithology and Physical Properties

Sample Alteration and Structural Measurements

Sample Location Map – Geology

Sample Location Map - Landsat

All samples were analysed at NTEL in Darwin using techniques outlined in **Cameco Standard Outcrop Sampling and Processing Procedures**.

Cameco Standard Outcrop Sampling and Processing Procedures

Sample G400 Geochemistry

All outcrop samples were processed using PIMA and TSG, with results in the following tables and figures.

Sample TSA Analysis (Identified Mineral Species)

Sample PIMA TSA Clay Distribution Map

The sandstone within the relinquished area is notably flat lying and well bedded, consisting of multiple 10-20cm thick units with occasional interspersed more massive units. Well-defined ripple marks and / or beds were noted throughout. Bedding dips rarely exceeded 5°. Pebbles and pebble units are rare, as are mud clasts. This sandstone has been assigned to the Marlgowa Sandstone Member of the Kombolgie Subgroup.

Clay determination analysis of PIMA Infrared spectra indicates the background sandstone is illite dominant.

Rare tight / closed 1-2mm thin quartz veins were noted throughout the area with no associated alteration, though ferruginous jointing was common. The average radiometric count for the relinquished area sandstone is ~50 – 80 cps.

Samples LP03C10105 and LP03C10106, mapped as brecciated Marlgowa Sandstone, may represent Cretaceous sediments.

CONCLUSIONS/DISCUSSION

Radiometric surveying and sandstone sampling located no notable anomalies. Any elevated radiometric and assay values are attributed to element scavenging by lateritisation of the ferruginous units and concentration along fractures and joint planes in the underlying Marlgowa Sandstone Member by groundwater flow.

Illite, dickite and kaolinite clay mineral assemblages dominate the samples analysed. This is consistent with regional clays in the Gumarrirnbang and Marlgowa Sandstone Members.

No further work is warranted or recommended within the area of relinquishment.

BIBLIOGRAPHY

Carson, L.J., Haines, P.W., Brakel, A., Pietsch, B.A., and Ferenczi, P.A. 1999. Milingimbi, Northern Territory. 1:25000 Geological Map Series. Northern Territory Geological Survey and Australian Geological Survey Organisation, Explanatory Notes SD 53-2.

Needham, R.S. 1988, Geology of the Alligator Rivers Uranium Field. Bureau of Mineral Resources Bulletin 224.

Sweet, I.P., Brakel, A.T., Carson, L., 1999, The Kombolgie Subgroup- a new look at an old 'formation'. AGSO Research Newsletter, 30; pages 26-28.