

FINAL RELINQUISHMENT REPORT FOR EL 8558

for the period
21/08/2003 to 24/12/2007

NORTHERN TERRITORY

Volume **1** of **1**

1:250,000 SHEET: Tanami SE52-15

1:100,000 SHEET: Tanami 4858

AUTHOR: F. Parker

TENEMENT HOLDERS: Newmont Tanami Pty Ltd

DISTRIBUTION: ☐ Northern Territory Department of Primary Industry, Fisheries and Mines
☐ Newmont Asia Pacific

The contents of this report remain the property of Newmont Tanami Pty Ltd and may not be published in whole or in part nor used in a company prospectus without the written consent of the Company.

SUMMARY

This is the final relinquishment report for EL 8558. As such, it details all exploration activity conducted over the licence for the period 21st August 2003 to 24th December 2007.

The tenement is located approximately 650km northwest of Alice Springs. The tenement is located approximately 12 km north of The Tanami Mine.

TABLE OF CONTENTS

1	INTRODUCTION	1
2	LICENCE DETAILS	1
2.1	LOCATION, ACCESS & PHYSIOGRAPHY	1
3	GEOLOGY	1
4	EXPLORATION HISTORY	2
5	REFERENCE LIST/ANNUAL REPORT BIBLIOGRAPHY	3

LIST OF TABLES

TABLE 1: Tenement Summary for EL 8558	1
---	---

LIST OF FIGURES

Figure No.	Title	Scale
1	Tenement Location and Access	1:500,000
2	Tenement Relinquishment	1:250,000
3	Exploration Index	1:500,000
4	Surface Geochemistry	1:250,000

LIST OF APPENDICES

APPENDIX 1:	DIGITAL SAMPLE & DRILLHOLE DATA
	EL8558_NTSG3SSASS_2007S.txt EL8558_WASL3SSAMP_2007S.txt

1 INTRODUCTION

This document is the final relinquishment report for EL 8558. It describes exploration activities over the tenement from 21st August 2003 to the 24th December 2007.

2 LICENCE DETAILS

Newmont Tanami Pty Ltd (Newmont) held and managed EL 8558.

TABLE 1: Tenement Summary for EL 8558

Licence	Grant	Surrender	Blocks	Km ²	Title Holder
EL 8558	21/08/2003	24/12/2007	112	360	100% Newmont Tanami Pty Ltd

2.1 LOCATION, ACCESS & PHYSIOGRAPHY

The tenement comprises 112 blocks situated approximately 12 km north of The Tanami Mine.

Main access is via the Tanami Track.

3 GEOLOGY

The Granites-Tanami Goldfields lie in the eastern part of the Early Proterozoic Granites-Tanami Inlier, which is part of the Northern Australian Orogenic Province (Plumb, 1990). The Inlier abuts the Arunta Complex to the south and east and is probably a continuation of the Halls Creek Orogen in Western Australia (Hendrickx, et al, 2000). The Inlier underlies younger cover sequences including the extensive Paleozoic Wiso Basin on its northeastern margin, and Victoria River Basin to the north. To the west, clastic sediments of the Middle Proterozoic Birrindudu Basin overlie and separate the Inlier from the similar age rocks in the Halls Creek Province.

The oldest rocks of the Tanami region belong to the Billabong Complex, a suite of Archaean age gneiss and schist. This is unconformably overlain by the Proterozoic MacFarlanes Peak Group (mafic volcanic and volcanoclastic rocks), followed by a thick succession of clastic sediments of the Tanami Group. (Hendrickx et al, 2000). A suite of syn- to post-deformation dolerites and gabbros are found intruding both the MacFarlane Peak and Tanami Groups.

Complex, polyphase deformation during the Barramundi Orogeny (1845 – 1840Ma) has affected the entire Granites-Tanami Inlier. It appears to have been largely controlled by two sets of regional scale fundamental crustal fractures that trend NNE and WNW. This is evidenced by the orientation of successive phases of macroscopic folding in the region and the consistent sympathetic trends of late tectonic faults.

Peak metamorphism during the Barramundi Orogeny reached amphibolite facies (The Granites Gold Mine), but is more generally greenschist facies through the Inlier (Callie Gold Mine). Contact metamorphic aureoles, commonly identified in pelitic schist units by randomly orientated andalusite porphyroblasts, are well developed at the margins of the syn- and post-orogenic granite plutons.

Localised extension followed, forming small basins which filled with shallow marine sediments to the west (Pargee Sandstone) and pillow basalts and turbiditic sediments to the east (Mt. Charles Formation).

Following the period of extension, widespread granite intrusion and volcanism followed in the period 1830 – 1810 Ma. At least three suites of granitic intrusives and two volcanic complexes are present. The last intrusion of (undeformed) granite occurred at around 1800 – 1795Ma, with intrusion of The Granites Suite (Hendrickx et al, 2000).

Residual hills of gently folded Carpentarian Gardiner Sandstone unconformably overlie Early Proterozoic lithologies. Younger flatlying Cambrian Antrim Plateau Basalts are also preserved as platform cover in areas protected from erosional stripping.

Tertiary drainage channels, now completely filled with alluvial and lacustrine clays and calcrete are a major feature of the region. Some drainage profiles are 10 km wide and greater than 100m deep.

A desert terrain comprising transported and residual colluvial cover sediments and aeolian sand blanket a large portion of the Inlier, with an estimated outcrop exposure of less than 10% of the early Proterozoic lithological units.

Gold mineralisation within the Newmont Tanami tenement holdings is dominantly hosted by the Tanami Group, a sequence of fine to medium-grained turbiditic metagreywackes with lesser amounts of metapelite, carbonaceous siltstone and schist, banded iron-formation, chert and calcsilicates. (Hendrickx et al, 2000). Owing to their more resistant nature, only the cherts and iron-formations and associated interbedded graphitic schists tend to outcrop above the sand plain. The interlayered pillow basalts and sediments of the Mt.Charles Formation at the Tanami Mine deposits also host significant gold mineralisation.

4 EXPLORATION HISTORY

During 2004 Newmont Exploration completed a comprehensive first pass surface sampling programme on a 800m x 800m and 800m x 400m grid.(412 samples). Nine rock chip samples were also taken.



5 REFERENCE LIST/ANNUAL REPORT BIBLIOGRAPHY

References

- Blake, D., Hodgson, I.M., and Muhling, P.C.**, 1979. Geology of The Granites-Tanami Region, Northern Territory and Western Australia, *Bur. Miner. Resour. Geol. Geophys. Aust. Bull.* 197.
- Davidson, A.A.** 1905. Journal of Explorations in Central Australia, by the Central Australian Exploration Syndicate, Limited, *South Australia Parliamentary Paper* 27.
- Gee, L.C.E.** 1911. General Report on Tanami Goldfield and District (Northwestern Central Australia). *South Australia Parliamentary Paper* 31.
- Hendrickx M.A., Slater K.R., Crispe A.J., Dean A.A., Vandenberg L.C., and Smith J.B.**, 2000. Palaeoproterozoic stratigraphy of the Tanami Region: regional correlations and relation to mineralisation – preliminary results. *Northern Territory Geological Survey. Geological Survey Record GS 2000-13.*
- Hodgson, C. J.**, 1975, Tanami, Northern Territory, 1:250,000 *Geological Series: Explanatory Notes.*
- Hossfeld, P.S.** 1940b. The Gold Deposits of The Granites-Tanami District, Central Australia. *Aer. Geol. Geophys. Surv. N.Aust., Northern Territory Report* 43.
- Mayer, T.E.** 1990. The Granites Gold Field, in *Geology of the Mineral Deposits of Australia and Papua New Guinea* (Ed F.E. Hughes) pp 719-724 (The Australasian Institute of Mining and Metallurgy: Melbourne).
- O'Driscoll, E.S.T.** 1990. Lineament Tectonics of Australian Ore Deposits, in *Geology of the Mineral Deposits of Australia and Papua New Guinea* (Ed F.E. Hughes) pp 33-41 (The Australasian Institute of Mining and Metallurgy: Melbourne).
- Plumb, K.A.** 1990. Halls Creek Province and The Granites-Tanami Inlier - regional geology and mineralisation, in *Geology of the Mineral Deposits of Australia and Papua New Guinea* (Ed F.E. Hughes) pp 681-695 (The Australasian Institute of Mining and Metallurgy: Melbourne).
- Tunks, A. J.**, 1996, Geology of the Tanami Gold Mine, Northern Territory, *PhD Thesis, University of Tasmania, Hobart.*

EL 8558 Reports to NT Department of Primary Industries Fisheries and Mines

- Muir, M.**, 2004. First annual report for EL 8558 Coomarie 21 August 2003 to 20 August 2004. Newmont CR 31690
- Muir, M.**, 2007. Interim report for EL 8558 Coomarie 21 August 2004 to 31 December 2004.. Newmont CR 31798



0 10 20
Kilometers

NEWMONT **NEWMONT EXPLORATION PTY LTD**

Tanami Project

EL 8558

LOCATION AND ACCESS

FIGURE 1



Author: F. Parker

Scale: 1:500 000

Drawn: V. Preedy

Date: Apr 2008

File: TAN_Lnd_Ten_A4_EL8558surr1.mxd Projection: Lat/Long (GDA 94)
\\AUS\NT\Tanami_Tenements\MXD\Reports\



Area Surrendered - 112 blocks



0 5 10
Kilometers

NEWMONT **NEWMONT EXPLORATION PTY LTD**



Tanami Project

EL 8558

TENEMENT RELINQUISHMENT

FIGURE 2

Author: F. Parker	Scale: 1:250 000
Drawn: V. Preedy	Date: Apr 2008
File: TAN_Lnd_Ten_A4_EL8558surr2.mxd Projection: Lat/Long (GDA 94) \\AUS\INT\Tanami_Tenements\MXD\Reports\	



▲ Surface Sampling



0 10 20

Kilometers

NEWMONT **NEWMONT EXPLORATION PTY LTD**



Tanami Project

EL 8558

EXPLORATION INDEX

FIGURE 3

Author: F. Parker	Scale: 1:500 000
Drawn: V. Preedy	Date: Apr 2008
File: TAN_Lnd_Ten_A4_EL8558surr3.mxd Projection: Lat/Long (GDA 94)	
\\AUS\NT\Tanami_Tenements\MXD\Reports\	



Surface Sampling

- ▲ Soils
- ▲ Rockchip



0 5 10
Kilometers

NEWMONT **NEWMONT EXPLORATION PTY LTD**

Tanami Project

EL 8558

SURFACE GEOCHEMISTRY

FIGURE 4

Author: F. Parker	Scale: 1:250 000
Drawn: V. Preedy	Date: Apr 2008
File: TAN_Lnd_Ten_A4_EL8558surr4.mxd Projection: Lat/Long (GDA 94) \\AUS\INT\Tanami\Tenements\MXD\Reports\	