

DRILLHOLE SUMMARY INDD001

HOLE: **INDD001** PROJECT: Indiana
DATE DRILLED: 9-15 June 2008 DRILL BY: Prime Drilling
GEOLOGIST: Michael Green
TARGET: 100 seimens ground EM conductor at 147 m.

MGA_E: 588484 MGA_N: 7445554
DIP: -60° DIP AZIMUTH: 260°true (255°mag)

HOLE DEPTH: 183.3 m
PRECOLLAR DEPTH: 7 m
ROD SIZE & DEPTH: HQ₃ 7.0-29.7 m; NQ₂ 29.7-183.3 m
DEPTH OF TRANSPORTED COVER: 4 m
DEPTH TO BASE OF OXIDATION: 23.1 m
WATER IN HOLE: no

COMMENTS: 50 mm casing to bottom of hole; 30 m steel HQ rods stuck at top of hole.

CORE MARKED & MEASURED: Yes
GEOPHYSICAL LOGS: mag sus & geiger on whole core
PHOTOGRAPHY: yes
CORE CUT: only mineralised intervals
SAMPLING: only mineralised intervals

SURVEYS

Depth	Dip	Mag Azimuth	True Azimuth		Depth	Dip	Mag Azimuth	True Azimuth
0	-60	255	260					
51	-61.8	252.2	257.2					
99	-62.9	252.0	257.0					
150	-63.8	251.8	256.8					

ORIENTATION MARKS: Chinograph spear at 51 and 102 m.

GEOLOGY/MINERALISATION:

Top 4 m comprises aeolian-colluvial sand and gravel. Bedrock geology comprises quartz-biotite-feldspar gneiss with lesser garnet-chlorite-kyanite and layered quartz-epidote-hornblende-calcite calc-silicate units. All of these units are interpreted to be metamorphosed sediments indicating an original package of interbedded siltstone, mudstone and carbonates.

Stringer to blebby pyrrhotite-pyrite-chalcopyrite is best developed in and around a dark green, foliated hornblende-chlorite-rich unit between 149.8-160.4 m. The chalcopyrite content is up to 2 % between 155.0-155.5 m, but is typically <1 %. The main host unit was initially interpreted as a metamorphosed mafic magmatic unit, but similar units in INDD002 are more consistent with a sedimentary origin, which is now the preferred interpretation. Sulphide stringers are folded in adjacent psammopelitic gneiss (eg 145.5 and 163 m) and mineralisation is interpreted to be pre-metamorphic. Some unusual units around this main

DRILLHOLE SUMMARY INDD001

mineralised zone are probably metamorphosed alteration associated with mineralisation. Mineralisation is also associated with magnetite.

From	To	Colour	Weather	Lithology	Variant	Mineral	Struct	α	Comments
0	4	rb	transported	aeolian/colluvium	sand/gravel				percussion precollar
4	7	gy	moderate	psammopelitic gneiss					percussion precollar
7	23.1	gy	slight	pelitic gneiss	medium- to coarse-grained foliated	bi qt fd cl ab	S0, S1	60	banded and layered; 20 cm wide calc-silicate ep-qt-ca-dp ?boudins - coarse-grained, more massive, probably elliptical
23.1	41.6	gy	fresh rock	psammopelitic gneiss	medium- to coarse-grained foliated	bi qt fd cl ab	S0, S1	60	40% interlayered calc-silicate - ep-qt-ca-dp - layers are generally banded subparallel to gneissosity in psammopelite to 3m wide; minor po>py
41.6	44.2	dgy	fresh rock	pelitic gneiss	coarse-grained	bi qt fd gt ky st py	S0, S1	60	
44.2	60.4	gy	fresh rock	psammopelitic gneiss	medium- to coarse-grained foliated	bi qt fd cl ab	S0, S1	60	50% interlayered calc-silicate ep-qt-ca-dp-ac; as for 23.1-41.6m
60.4	64.6	dgy	fresh rock	pelitic gneiss	coarse-grained foliated	bi qt fd cl ab			minor calc-silicate; possible large antiform - axial plane subparallel to foliation
64.6	99.5	lgy	fresh rock	calcsilicate	medium-grained foliated	ep cl ac ab ca	S0, S1	70	minor psammopelitic to pelitic layers; rootless folds; large fold at 90m; trace po-py
99.5	109.5	gy	fresh rock	pelitic gneiss	coarse-grained foliated	bi qt ft fd	S0, S1	70	minor pelite and calc-silicate
109.5	113.9	gg	fresh rock	calcsilicate	medium-grained foliated	ep qt ca ac	S0, S1	70	minor psammopelite to pelite; <<1% disseminated py
113.9	125.7	gy	fresh rock	psammopelitic gneiss	medium-grained foliated	bi qt fd gt	S0, S1	60	minor pelite and calc-silicate; <1% disseminated py & stringer py-po
125.7	128.4	gr	fresh rock	calcsilicate	medium-grained, massive	qt ep dp ca			
128.4	129.25	gy	fresh rock	psammopelitic gneiss	medium-grained foliated	bi qt fd	S0, S1	60	layered; minor py bands subparallel to foliation
129.25	131.1	wh	fresh rock	calcsilicate	mg ma	ca ep dp			<<1% disseminated py
131.1	139.4	gy	fresh rock	psammopelitic gneiss	medium- to coarse-grained foliated	bi qt fd cl			minor calc-silicate layers; <<1% py bands subparallel to foliation
139.4	149.8	gy wh	fresh rock	calcsilicate	coarse-grained, massive	qt ca bi pl dp hb cl			includes qt-rich domain, dalmation rock with coarse-grained hb-cl-pl-qt & interconnected semi-massive py-po-mt-cp; metamorphosed alteration; folded pelite 145-146m
149.8	160.4	dgr	fresh rock	calcsilicate	medium-grained	cl hb ab py po mt			weakly foliated massive-blebby po>py>>cp; minor coarse-grained mt; possible pyrobitumen with \$\$
160.4	164.6	gy	fresh rock	pelitic gneiss	coarse-grained foliated	bi ky qt cl po py mt	S0, S1	60	py-po in foliation and folded; some py-po becoming stringer
164.6	175	gg	fresh rock	calcsilicate	medium-grained foliated	qt hb ep ca	S0, S1	60	some large fold hinges
175	183.3	gy	fresh rock	psammopelitic gneiss	medium-grained foliated	bi qt cl fd	S0, S1	60	minor calc-silicate bands; folds

