

System	Stage/ Substage	S. Africa	Australia				
		Karoo Basin (Anderson, 1977)	Collie Basin, W. Australia (Backhouse, 1991)	E. Australia (Price <i>et al.</i> , 1985)		E. Australia (Kemp <i>et al.</i> , 1977)	Canning Basin, W. Australia (Kemp <i>et al.</i> , 1977)
Permian (part)	? Ufimian		<i>D. parvithola</i>	PP5	PP5.1	upper Stage 5	Unit VII
			<i>P. rugatus</i>	PP4	PP4.3	lower Stage 5c	
		4d	<i>D. ericianus</i>		PP4.2	lower Stage 5b	
	4c	<i>D. granulata</i>	PP4.1		lower Stage 5a		
	4b						
	4a		<i>M. villosa</i>	PP3.3	upper Stage 4b		
	3d						
	3c	(consistent occurrence of <i>P. sinuosus</i> )				PP3	PP3.2
	3b		PP3.1	lower Stage 4	Unit V		
	Aktastinian		3a	<i>M. trisina</i>	PP2		PP2.2
		2d	PP2.1			Stage 3a	
		2c					
	Sterlitamakian Tastubian	2b 2a	<i>P. pseudoreticulata</i>				Unit III
		? Asselian	1	<i>P. confluens</i>	PP1	Stage 2	Unit II
	Stage 2						

Figure 5.3 Inferred correlation of Permian palynozonations established in Australia and the Karoo Basin of South Africa. Adapted from Backhouse (1991, fig. 10).